

CANVIL-5%

**FUNGICIDE
SUSPENSION CONCENTRATE**

VAPCO

Class:
Triazole

A systemic fungicide effective as both protective and curative action against a broad range of fungal diseases such as Scab, Powdery mildew, Rust, Rots on a wide range of agricultural crops.

COMPOSITION: Each liter contains:

HEXAICONAZOLE

5 % W/V (A.I)

PROPERTIES:

CANVIL-5%: Systemic, protective and curative fungicide to control a broad range of fungal diseases on a wide range of agricultural crops.

CANVIL-5%: Slightly toxic to mammals and not phytotoxic if used as recommended.

CANVIL-5%: Controls Powdery mildew, Rust, Brown rot, Black rot and other fungal diseases.

CANVIL-5%: Slightly toxic to bees and toxic to fish.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE ML / 20 LT WATER	SAFETY PERIOD (DAYS)
Cucumber, Tomato, Squash, Pepper	Powdery mildew	4 - 8	14
Cabbage, Cauliflower	Powdery mildew	10 - 20	7
Onion, Garlic, Peas	Powdery mildew	4 - 8	14
Carrot	Powdery mildew	10 - 20	7
Leek	Powdery mildew	10 - 20	14
Watermelon, Muskmelon	Powdery mildew	4 – 8	4
Plum, Apricots	Brown rot, Powdery mildew, Rust	8 – 20	14
Apple, Mango	Scab, Powdery mildew, Rust	4 - 20	14
Grapes	Powdery mildew, Black rot	4 – 8	14
Strawberry	Powdery mildew	6 - 10	2
Ornamentals	Powdery mildew, Rust	4 – 8	---

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY:

Compatible with most pesticides, except alkaline compounds, it's favorable to make compatibility test before use.

*** OTHER FORMULATIONS & TRADE NAME:**

CANVIL – 10% EC, CANVIL – 10% SC, VAPCOVIL – 5% SC.

• For more details about first aid & precautions please refer to first aid & precautions index.