

CARTAP-50%

**INSECTICIDE
SOLUBLE POWDER**

VAPCO

Class:
**2-dimethylaminopropane
-1,3- dithiol**

Systemic insecticide used to control chewing and sucking insects in all developing stage on many agricultural crops.

COMPOSITION: Each kg contains:

CARTAP

50% W/W (A.I)

PROPERTIES:

CARTAP-50%: Systemic insecticide acts by contact and stomach action.

CARTAP-50%: Used to control many chewing and sucking insects on a wide range of agricultural crops.

CARTAP-50%: Moderately toxic to mammals, toxic to fish and low toxic to bees.

CARTAP-50%: There is no residue in animal tissue.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE GM / 20 LT. WATER	SAFETY PERIOD (DAYS)
Cabbage	Diamond-back moth	25	14
Cucurbits	Thrips, Aphids, Leafworm, Flea Beetle, Squash Beetle	19 - 25	10
potato	Potato Tuber moth	25	7
Broccoli	Cutworm, Aphids	25	7
Spanish	Thrips	25	7
Garlic	Thrips, Cutworm	19 - 25	10
Onion	Thrips, Cutworm	19 - 25	10
Peas	Leafminer	25 – 50	1
Corn	European corn borer	20	21
Peach	Cutworm, Aphids	19 - 37.5	10
Mango	Leafhoppers, Twig borer, Tip borer	19 - 25	10
Rice	Whorl Maggot, Rice stemborer, Green leafhopper, Brown planthopper	37.5 - 62.5	10
Ornamental	Thrips	25	5

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

48 hours after the last application.

COMPATIBILITY:

Compatible with most pesticides except alkaline materials.

*** OTHER TRADE NAME: VARTAP – 50% SP.**

• For more details about first aid & precautions please refer to first aid & precautions index.