

CHLORONIL-75%

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Chloronitrile

Non-systemic fungicide, protective and curative fungicide to control many fungal diseases.

COMPOSITION: Each kg contains:

CHLOROTHALONIL

75 % W/W (A.I)

PROPERTIES:

CHLORONIL-75%: Non-systemic fungicide, an excellent protection Fungicide against many fungal diseases alone or as a part of controlling programme.

CHLORONIL-75%: Multi-site activity on disease resistance.

CHLORONIL-75%: Rain fast and weather proof; remains, effective after rainfall and/or irrigated crops.

CHLORONIL-75%: Non-phytotoxic when used as recommended.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE GM/ 20 LT. WATER	SAFETY PERIOD (DAYS)
Potatoes	Late blight, Early blight, Root rot, Wilt (<i>Rhizoctonia</i>).	30 – 40	7
Cucumber, Squash	Powdery mildew, Sclerotinia, Wilt (<i>Rhizoctonia</i>).	30 – 40	7
Tomatoes	Late blight, Anthracnose, Septoria spot, Brown rot, Sclerotinia, Wilt (<i>Rhizoctonia</i>).	30 – 40	7
Pepper, Eggplant	Alternaria Spot, Anthracnose, Brown Rot, Downy mildew, Septoria Spot	30 – 40	3
Cabbage, Cauliflower, Broccoli	Alternaria Spot, Downy mildew	30 – 40	10
Onion	Gray Mold, Downy mildew	30 – 40	10
Muskmelon	Anthracnose, Downy mildew, Black Root Rot	40	3
Apple	Brown Rot ,Canker	30 – 40	15
Peach, Apricot	Brown Rot ,Canker	30 – 40	15
Vines	Downy mildew, Gray mold, Fire blight, Powdery mildew	30 – 40	15
Bean	Anthracnose	26	30

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE GM/ 20 LT. WATER	SAFETY PERIOD (DAYS)
Peas	Anthrachnose, Gray Mold	10	3
Cereals	Rust, Powdery mildew, Spots	2 – 3 kg / ha	7
Ornamental, Roses and Turf	Powdery mildew, Brown rot	35 – 50	7

Hectare = 10000 m²

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

NOTES:

- Can be used at rate 50 gm / 20 lt. water as irrigation to control soil fungal diseases as wilt (*Rhizoctonia*).
- To obtain best results, it is advisable to spray at the beginning of season as protective spray. don't wait until symptoms of disease appear.
- Repeat spraying every 7-10 days.
- At severe infection repeat spraying at shorter interval and increase the dosage.

RE-ENTRY PERIOD:

48 hours after the last application.

COMPATIBILITY:

Compatible with most common pesticides, it's advisable to make mixing trial before mixing the product and it is not advisable to mix with foliar fertilizers.

*** OTHER FORMULATIONS: CHLORONIL – 75% SC**

• For more details about first aid & precautions please refer to first aid & precautions index.