

# COCCIDOLE

Powder



**COMPOSITION:** Each gm contains:

**Clopidol 125 mg**

**PROPERTIES:**

- Clopidol is of pyridinol group which acts on first generation schizonts of coccidia , and thus prevents the disease occurrence.
- It dose not develop any immunity against coccidiosis. It is non-toxic for non-host species, very safe to be used and is compatible with other medicaments including antibiotics, so it can be administered with other medicaments except other anticoccidials.
- It helps and improves metabolism, so it promotes growth and improves performance, feed conversion and weight gain.

**INDICATIONS:**

For prevention and treatment of most species of Eimeria strains causing caecal and intestinal coccidiosis in poultry, rabbits, turkeys and game birds ( partridges, pheasants, guinea-fowls).

**DOSAGE:**

- **Prevention:** Start medication at day old of the flock.
- **Poultry:** 1 kg per ton of feed continuously up to 16 weeks old.
- **Rabbits:** 1.6 kg per ton of feed continuously.
- **Turkeys:** 800 gm per ton of feed continuously up to 12 weeks old.
- **Game birds:** 1 kg per ton of feed continuously.
- **Treatment:** According to the severity of the case, double the preventive dose up to 2-4 times for 2-4 days, then back to the preventive dose.

**REMARK:**

- Not recommended for layers.
- Not mixed with other anticoccidials.

**SAFETY PERIOD:** For meat consumption : 7 days.

**STORAGE:** Keep in a dry, dark place below 30 °C.

**PACKING:** (1, 5, 10, 25) kg.