

# COMMANDO-35%

INSECTICIDE  
SUSPENSION CONCENTRATE

VAPCO

Class:  
Neonicotinoid

A systemic insecticide, with a broad spectrum efficiency especially against sucking insects such As Aphids, Leafhoppers, Thrips, Beetles And Leafminer On Tomato, Pepper, Cabbage, Lettuce, Cauliflower, Eggplant, Beans, Carrots, Apple, Pear, Plum, Peach, Apricot, Nectarine, Tobacco and Ornamentals. Also it is used to control all species of subterranean Termites in the field.

**COMPOSITION:** Each liter contains:

**IMIDACLOPRID**

**35% W/V (A.I)**

## PROPERTIES:

**COMMANDO-35%:** A systemic insecticide, with a broad spectrum efficiency especially against sucking insects such as Aphids, Hoppers, Thrips, Beetles, and Leafminer on Tomato, Pepper, Cabbage, Lettuce, Cauliflower, Eggplant, Beans, Carrots, Apple, Pear, Plum, Peach, Apricot, Nectarine, Tobacco, and Ornamentals.

**COMMANDO-35%:** Highly efficient in controlling species of subterranean termites in the field.

**COMMANDO-35%:** Used as contact and stomach action.

**COMMANDO-35%:** Controls a wide range of pests such as Aphids, Leafhopper, Thrips, Whitefly, Wireworms, Colorado Beetles, Wooly Aphid, Scale Insects.

**COMMANDO-35%:** Has a long lasting control.

**COMMANDO-35%:** Controls insects which became resistant against other pesticides.

## USES AND APPLICATION RATE:

### 1- TERMITE CONTROL IN THE FIELD:

**COMMANDO-35% SC** is used to control subterranean Termites in vegetable fields at the rate of 25-50 ml/20 lt. of water applied as follow:

- Complete coverage through spraying over the soil before planting, ploughing and mixing of the sprayed soil to the depth of 5-10 cm.
- Spray over planting lines after seeding, then complete coverage with soil over the treated seeds.

### 2- AGRICULTURAL USES:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Citrus	Psylla, Aphid, Scale Insect, Mealybug	9 - 11 Irrigate before 2 weeks of flowering	30
Peaches, Nectarines, Plums, Apricots, Cherries, Almond, Apples, Pears	Aphid, Leafminer. Pear Psylla, Wooly Aphid	4-6 6-14	14
Mango	Leafhoppers, Scale insect	2-3	30

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Grapes	Aphid, Leafhoppers, Mealybug, Thrips.	1 - 9	30
Strawberries	Aphid	2 - 29	14
Cucumber, Squash, Pepper, Tomatoes.	Aphids, Leafhoppers.	3 - 6	7
Cabbage, Lettuce, Cauliflower, Eggplant, Beans, Carrots	Thrips, Leafminer, Whiteflies.	6 - 14	
Potatoes	Colorado Beetle. Aphid, Leafhopper, Cutworms.	6 - 14 3 - 6	30
Tobacco	Aphid, Whitefly	3 - 6	21
Okra	Aphid, Thrips	1	21
Soybeans	Aphid	2 - 6	30
Sugar Beet	Aphids, Leafhoppers, Whitefly, Flea Beetles	5 - 11	30
Sunflower, Corn	Aphid, Whitefly	6	21
Rice	Leafhoppers, Leaf Beetles, Rice Stink Bug, Rice Weevil, Thrips.	1 - 6	37
Cotton	Aphids, Bugs, Leafhoppers, Thrips, Whitefly, Beetles	1 - 20	28
Coffee	Leafminer, Scale insect, Bugs.	1 - 9	45
Tea	Leafhoppers, Thrips, Aphids, Leafrollers.	2 - 17	14
Palm	Scale Insect, Leafhopper, Beetles.	9 - 11 ML/ 5 Liter water for each tree yearly.	After harvesting
Roses	Aphid	3	---
Ornamentals	Aphid, Thrips, Whitefly, Beetles	6 - 12	---

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest

**RE-ENTRY PERIOD:**

48 hours after the last application.

**COMPATIBILITY:**

Compatible with most pesticides except alkaline compounds, it's preferable to use the product immediately after mixing.

• For more details about first aid & precautions please refer to first aid & precautions index.