

COMMANDO-5%

INSECTICIDE
GRANULES

VAPCO

Class:
Neonicotinoid

Systemic insecticide, with broad-spectrum efficiency against many insect pests such as Aphids, Leafhoppers, Thrips, Beetles, Leafminer and other sucking and chewing insects on a wide range of agricultural crops.

COMPOSITION: Each kg contains:

IMIDACLOPRID

5% W/W (A.I)

PROPERTIES:

COMMANDO-5%: Systemic insecticide with broad spectrum efficiency against sucking insects and some chewing insects on vegetables, fruit trees, and ornamentals.

COMMANDO-5%: Has a contact and stomach action.

COMMANDO-5%: Control a wide range of pests such as Aphids, Leafhoppers, Thrips, Whitefly, Wireworms, Colorado beetle, Wooly aphids, Scale insects ... etc.

COMMANDO-5%: Has a long lasting effect up to 70 days.

COMMANDO-5%: Controls insects which became resistant to other pesticides.

COMMANDO-5%: Moderately toxic to mammals, toxic to bees, fish and birds.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE	SAFETY PERIOD (DAYS)
Tomato, Pepper, Lettuce, Cabbage, Cauliflower, Eggplant, Bean, Carrots, Squash, Pumpkin, Watermelon, Muskmelon, Cucumber.	Aphids, Leafhopper	0.1 – 0.2 gm / m line	7
	Thrips, Whitefly, Leafminer	0.3 – 0.5 gm / m line	
Potato	Colorado Beetles, Leafhopper, Aphid Wireworm	0.5 – 1.0 gm / m line	30
		1.0 – 1.5 gm / m line	
Apple, Pear, Quince, Plum, Peach, Almond, Apricots, Cherries, Nectarine	Aphids, Leafminers.	20 – 40 gm / tree	14
	Pear Psylla, Wooly Aphid.	30 – 60 gm / tree	
Lemon, Orange, Pummelo, Mandarin, Grapefruit	Aphid, Psylla	20 – 40 gm / tree	14
	Thrips, Scale Insects	30 – 60 gm / tree	

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE	SAFETY PERIOD (DAYS)
Date Palm	Dubas Bug, Lesser Date Moth.	20 – 40 gm / tree	30
Rice	Brown Leafhopper, Green Leafhopper, Small Brown Leafhopper, White Back Leafhopper	10 - 20 gm/ seedbed	---
	Rice Leaf Beetle		
Tobacco	Aphids. Whitefly.	0.1 – 0.3 gm / plant 0.4 – 0.6 gm / plant	21
Ornamentals	Aphids, Thrips, Whitefly, Beetles.	20 – 50 gm / m ²	---

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

12 hours after the last application.

COMPATIBILITY:

Compatible with most pesticides except alkaline materials.

• For more details about first aid & precautions please refer to first aid & precautions index.