

COPROXIDE

**FUNGICIDE
WETTABLE POWDER**

VAPCO

Class:
Inorganic compound

A protective fungicide and bactericide on a wide range of agricultural crops.

COMPOSITION: Each kg contains:

**COPPER HYDROXIDE
(EQV. 50% W/W COPPER (A.I))**

77% W/W

PROPERTIES:

COPROXIDE: Is a contact fungicide and bactericide with a preventive action against a wide range of diseases.

COPROXIDE: Adhere to the plant surface as the rainfall has low effect on washing it.

COPROXIDE: Control bacterial infections on agricultural crops such as Angular leaf spot.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE GM / 20 LT. WATER	SAFETY PERIOD (DAYS)
Peas	Powdery Mildew	32.5 - 75	7
Beans	Bacterial Spot, Rust	32.5 - 75	7
Cabbage	Black Rot, Black Leaf Spot	32.5 - 75	7
Cauliflower	Downy Mildew, Alternaria Leaf Spot	32.5 - 75	7
Broccoli	Downy Mildew, Leaf Spot, Black Mold	32.5 - 75	7
Carrots	Cercospora Leaf Spot	32.5 - 75	7
	Alternaria Leaf Spot	20	7
Celery	Early and Late Blight	32.5 - 75	7
Cucumber	Anthrachnose, Septoria Leaf Spot, Cercospora Leaf Spot, Fruit Rot, Downy Mildew	32.5 - 75	7
Watermelon	Anthrachnose, Cercospora Leaf Spot, Downy Mildew	32.5 - 75	7
Squash	Anthrachnose, Angular Spots, Downy Mildew	17 - 35	7
Potato	Early and Late Blight	10 - 45	15
Pepper	Bacterial Spot	32.5 - 75	7
Tomatoes	Bacterial Spot, Early and Late Blight, Anthracnose	20 - 45	10
Eggplant	Anthrachnose, Blight	20	10
Onion, Garlic	Purple Blotch, Downy Mildew	32.5 - 75	7
	Bacterial Spot	10 - 17	

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE GM / 20 LT. WATER	SAFETY PERIOD (DAYS)
Banana	Sigatoka Leaf Spot	32.5 - 75	7
Mango	Anthracnose	32.5 - 75	7
Grapes	Anthracnose, Mildews, Black Rot	32.5 - 75	7
Almond, Apricot, Plum, Peach, Cherry	Flower Rot, Shot Hole, Leaf Curl, Scab	35 - 50	15
Apple, Pear	Anthracnose, Black Spot, Fire Blight, Apple Scab	50 - 70	7
Orange, Lemon, Mandarin, Grapefruit	Black Rot, Spot	45 - 90	21
Olives	Peacock Spot, Olive knot	35 - 40	21
Cacao	Black Pod	32.5 - 75	21
Coffee	Leaf Rust, Coffee Berry Disease	32.5 - 75	21
Peanut	Cercospora Leaf Spot	32.5 - 75	21
Groundnut	Tikka Leaf Spot	75	---
Tea	Blister Blight	10	---
Alfalfa	Leaf Spot	25	10 – 14
Wheat, Barley	Leaf Blotch	15 - 25	10
Rice	Blast, Sheath Blight, Bacterial Leaf Strike	32.5 - 75	7

CAUTION:

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- Low mammalian toxicity, not phytotoxic for treated plants except some species of Apples (Red Apple)

NOTE:

Used inside plastic houses on Tomatoes, Pepper and Eggplant.

RE-ENTRY PERIOD:

48 hours after the last application.

COMPATIBILITY:

Compatible with most other pesticide except acidic compounds or Calcium sulphate.

*** OTHER TRADE NAME: VAPCOPRO - 77 % WP .**

• For more details about first aid & precautions please refer to first aid & precautions index.