

# CRIPTAN-50%

FUNGICIDE  
WETTABLE POWDER

VAPCO

Class:  
Phthalimide

Fungicide with protective and curative action controls a wide range of fungal diseases on a wide range of crops.

**COMPOSITION:** Each kg contains:

**CAPTAN**

**50% W/W (A.I)**

**PROPERTIES:**

**CRIPTAN-50%:** Used to control various fungal diseases affecting vegetation and seedlings.

**CRIPTAN-50%:** Control many of fungal diseases, affecting crop plants such as Spots, Leaf curl, Late and early blight and Seedlings diseases.

**CRIPTAN-50%:** With low mammalian toxicity.

**USES AND APPLICATION RATE:**

CROP	DISEASE	APPLICATION RATE GM / 20 LT. WATER	NUMBER OF APPLICATION	SAFETY PERIOD (DAYS)
Cucumber, Squash, Pumpkin	Anthrachnose, Scab	65 - 135	Apply every 5-7 days	2
Tomatoes	Anthrachnose, Early Blight, Late Blight, Leaf Spot, Septoria Leaf Spot	40 - 80	Apply every 7-10 days	2
Broccoli, Cabbage, Cauliflower, Eggplant, Pepper, Tomatoes, Rose.	Seedling Wilt, Root Rot	125 – 225 gm/ 100 m <sup>2</sup>	Work into top 8 - 10 cm of soil before planting.	---
Broad beans, Peas, Leek	Damping Off, Seedling Blight, Spots, Dieback,  Leaf Blight	40 - 50	Spraying or drenching  Spraying	10
Potatoes	Early Blight, Late Blight	40 - 50	Apply every 10-15 days	10
Celery	Leaf Spot, Late Blight	60 - 80	---	7
Garlic, Onions	Gray Mold, Purple Blotch, Downy Mildew	60 - 80	---	7

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE GM / 20 LT. WATER	NUMBER OF APPLICATION	SAFETY PERIOD (DAYS)
Carrots	Ring Spot, Cercospora Blight	60	As necessary	14
Lettuce	Leaf Spot	90	As necessary	7
Tobacco	Late Blight, Canker, <i>Fusarium</i> Wilt	70 - 80	---	7
Lemon, Orange, Pummelo, Mandarin, Grapefruit	Fruit Rot, Gummosis, Dieback	40 - 50	Spray two times "at the beginning of June & September"	14
Grapes	Dead Arm	40	Spraying	7
Apricot, Plum, Peach, Almond, Cherry	Brown Mold, Peach Leaf Curl	40 - 50	Spray before bud open	7
Apple, Pears	Black Spot, Leaf Spot, Sooty Blotch, Ripe Spot	20 - 40	Spraying	7
Avocado	Anthraxnose, Scab	70 - 80	---	7
Coffee	Coffee Berry Disease, Cercospora Blight, Brown Leaf Spot, Damping Off, Black Root Rot	60 - 100	---	7
Pineapple	Root Rot, Bud Rot	40 - 100	---	7
Strawberry	Gray Mold, Leaf Spot	90 - 135	Spray in the beginning of growth in spring and repeat the spraying every week	2
Ornamentals	Leaf Spot, Flower Blight, Bulb Mold	40 - 50	Spraying or seed treatment	---

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

**RE-ENTRY PERIOD:**

24 hours after the last application.

**COMPATIBILITY:**

Incompatible with all alkaline compounds, Oil sprays, TEPP compound, Emulsifiable Parathion compounds.

**\* OTHER FORMULATION: CRIPTAN - 80 % WP**

• For more details about first aid & precautions please refer to first aid & precautions index.