

# CURTINE-V

**FUNGICIDE  
WETTABLE POWDER**

**VAPCO**

Class:  
Dithiocarbamate<sup>(1)</sup>  
cyanoacetamide oxime<sup>(2)</sup>

**Effective fungicide to control a broad spectrum of fungal diseases such as Downy mildew and Blights.**

**COMPOSITION:** Each kg contains:

<b>MANCOZEB<sup>(1)</sup></b>	<b>46.5%W/W (A.I)</b>
<b>CYMOXANIL<sup>(2)</sup></b>	<b>4 %W/W (A.I)</b>

## **PROPERTIES:**

**CURTINE-V :** Highly absorbed by Plants, ensuring maximum utilization of the pesticide before being loosed by rain.

**CURTINE-V :** Protective Fungicide, prevent the formation of Mycelium at the incubation period and 4 days after infection.

**CURTINE-V :** Due to its dual structure its exhibits a tremendous efficiency against primitive fungi such as Downy mildew and blights on crops mentioned in the table of uses.

**CURTINE-V :** Could be sprayed three days after rain to maintain good results.

## **USES AND APPLICATION RATE:**

<b>CROP</b>	<b>DISEASE</b>	<b>APPLICATION RATE GM/ 20 LT. WATER</b>	<b>SAFETY PERIOD (DAYS)</b>
<b>Grapes.</b>	Downy mildew.	40 - 50	14
<b>Melons, Cucumbers</b>	Downy mildew.	40 - 60 Repeat 7 - 14 Days	---
<b>Potatoes, Tomatoes.</b>	Late blight, Downy mildew.	40 - 50	15
<b>Lettuce.</b>	Downy mildew.	40 - 50	15
<b>Chillies</b>	Late blight	40 - 60 Repeat 7 - 14 Days	---
<b>Onion, Garlic.</b>	Downy mildew.	40 - 50	14
<b>Tobacco.</b>	Downy mildew.	40 - 50	14

## **CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

## **RE-ENTRY PERIOD:**

24 hours after the last application.

## **COMPATIBILITY:**

Compatible with most other pesticides, compatibility test is recommended before use.

## **\* OTHER FORMULATIONS:**

**CURTINE-V - 72% WP (MANCOZEB 64% + CYMOXANIL 8%),**

**CURTINE-V - 76.42% WP (MANCOZEB 70.5% + CYMOXANIL 5.92%)**

• For more details about first aid & precautions please refer to first aid & precautions index.