

CYCO – S

Water Soluble Powder

VAPCO
ANTICOCCIDIAL

COMPOSITION: Each gm contains:

Sulphadimidine Sodium	320 mg
Diaveridine HCL	10 mg

PROPERTIES: An effective antioocidial and antibacterial product.

INDICATIONS:

Prevention and treatment of Salmonella and most species of Eimeria strains (Acervulina, Necartix, Tenella, Brunetti, Maxima, Hagani, Mitis, Praeacox) causing caecal, hepatic and intestinal coccidiosis in poultry (chickens, turkeys) and rabbits.

DOSAGE:

- **Poultry:** 100 gm per 200 litres of drinking water for 3 consecutive days, followed by plain water for 2 days, then apply medicated water for the next 2 days.
- **Rabbits:** 1 gm per 4 litres of drinking water for 3-5 days.
- **Via feed:** 2 kg per ton of feed for 3 days.
- **In severe cases:** The dose can be doubled and for the same period.

SAFETY PERIOD: For meat consumption: 7 days.

REMARKS:

- Used for broilers only.
- If used in laying hens, decrease in eggs production will be noticed.
- The use of Neomycin or Colistin - 600 for 3-5 days is highly recommended during coccidia treatment.
- Ensure adequate water intake during treatment to reduce the risk of crystalluria.

STORAGE: Keep in a dry, dark place below 30 °C

PACKING: (100, 250, 500, 1000) gm.