

DELTARIN-2.5%

**INSECTICIDE
EMULSIFIABLE CONCENTRATE**

VAPCO

**Class:
Pyrethroid**

One of the strongest insecticides in the pyrethroidal group which control a wide range of insects.

COMPOSITION: Each liter contains:

DELTAMETHRIN

2.5% W/V (A.I)

PROPERTIES:

DELTARIN - 2.5%: A non-systemic pyrethroidal insecticide with contact and stomach action.

DELTARIN - 2.5%: Quick action against a wide range of insect pests under different conditions.

DELTARIN - 2.5%: One of the strongest insecticides in the pyrethroidal group.

DELTARIN - 2.5%: Is not affected by light or temperature with a long activity period.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Tomato, Peppers Cucumber (Under Plastic House)	Whitefly, Tomato fruit worm, Leafminers, Thrips, Aphids, Lepidopterous larvae.	10 - 15	3
Cabbage, Cauliflower	Cabbage leaf worms, Lepidopterous larvae.	10 - 12	7
Potatoes	Leafroller, Aphids, Thrips, Colorado beetle larvae, Potato tuber moth	10 - 15	3
Onions	Onion fly, Thrips, Leafminers	10 - 15	7
Apples, pears, peach, Apricot	Caterpillars, Psylla, Leafminers, Codling moth, Pear sucker, Fruit fly	10 - 15	7
Mango	Hoppers, Tip and Twig borers	10 - 12.5	7-10 (do not spray at 29 - 35 days after bud break)
Banana	Flower thrips, Aphids	12.5	7 - 10
Legumes (Beans, Soybeans, etc.)	Bean fly, Flea beetles, Bugs, Thrips, Cutworms, Armyworms, Pod borers, Cutworms, Armyworms, Semi loopers, Bean lycaenid	10 - 12.5	For pre-flowering insect, 7 - 10 days are sufficient

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Legumes (Beans, Soybeans, etc.)	Pod borers, Cut worms, Armyworms, Semi loopers, Bean lycaenid	12.5	For post-flowering insect, use 12.5 ml/20L water at 30,40 & 50 days
Corn	Corn borer, Corn earworm, Seedling maggots	12.5	7 - 10
Rice	Whorl maggot, Green leafhopper, Stem borer, Rice bug, Rice black bug, Caseworm	10 - 12.5	7 - 14 days depending on the pest infestation level
Tobacco	Budworms, Armyworms, Cutworms	10 - 12.5	7 - 10
Cacao	Pod borer, Aphids	12.5	2 - 4

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY:

Compatible with most pesticides.

*** OTHER FORMULATION: DELTARIN - 0.05% (DUSTING POWDER).**

• For more details about first aid & precautions please refer to first aid & precautions index.