

# DELTARIN-2.8%

**INSECTICIDE  
EMULSIFIABLE CONCENTRATE**

**VAPCO**

**Class:  
Pyrethroid**

One of the strongest insecticides in the pyrethroidal group which control a wide range of insects.

**COMPOSITION:** Each liter contains:

**DELTAMETHRIN**

**2.8% W/V (A.I)**

**PROPERTIES:**

**DELTARIN - 2.8%:** A non-systemic pyrethroidal insecticide with contact and stomach action.

**DELTARIN - 2.8%:** Quick action against a wide range of insect pests under different conditions.

**DELTARIN - 2.8%:** One of the strongest insecticides in the pyrethroidal group.

**DELTARIN - 2.8%:** Is not affected by light or temperature with a long activity period.

**USES AND APPLICATION RATE:**

<b>CROP</b>	<b>PESTS</b>	<b>APPLICATION RATE ML / 20 LT. WATER</b>	<b>SAFETY PERIOD (DAYS)</b>
Tomato, Peppers Cucumber (Under Plastic House)	Whitefly, Tomato fruitworm, Leafminers, Thrips, Aphids, Lepidopterous larvae.	10 - 15	3
Cabbage, Cauliflower	Cabbage leafworms, Lepidopterous larvae.	10 - 12	7
Potatoes	Leaf roller, Aphids, Thrips, Colorado beetle larvae, Potato tuber moth	10 - 15	3
Onions	Onion fly, Thrips, Leafminers	10 - 15	7
Apples, pears, peach, Apricot	Caterpillars, Psylla, Leafminers, Codling moth, Pear sucker, Fruit fly	10 - 15	7
Mango	Hoppers, Tip and Twig borers	10 - 12.5	7-10 (do not spray at 29 - 35 days after bud break)
Banana	Flower thrips, Aphids	12.5	7 - 10
Legumes (Beans, Soybeans, etc.)	Bean fly, Flea beetles, Bugs, Thrips, Cutworms, Armyworms	10 - 12.5	For pre-flowering insect, 7 - 10 days are sufficient

• For more details about first aid & precautions please refer to first aid & precautions index.

<b>CROP</b>	<b>PESTS</b>	<b>APPLICATION RATE ML / 20 LT. WATER</b>	<b>SAFETY PERIOD (DAYS)</b>
<b>Legumes (Beans, Soybeans, etc.)</b>	Pod borers, Cutworms, Armyworms, Semi loopers, Bean lycaenid	12.5	For post- flowering insect, use 12.5 ml/20 L water at 30,40 & 50 days
<b>Corn</b>	Corn borer, Corn earworm, Seedling maggots	12.5	7 - 10
<b>Rice</b>	Whorl maggot, Green leafhopper, Stem borer, Rice bug, Rice black bug, Caseworm	10 - 12.5	7 - 14 days depending on the pest infestation level
<b>Tobacco</b>	Budworms, Armyworms, Cutworms	10 - 12.5	7 - 10
<b>Cacao</b>	Pod borer, Aphids	12.5	2 - 4

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

**RE-ENTRY PERIOD:**

24 hours after the last application.

**COMPATIBILITY:**

Compatible with most pesticides.

**\* OTHER FORMULATION: DELTARIN - 0.05% (DUSTING POWDER).**

• For more details about first aid & precautions please refer to first aid & precautions index.