

DEWORMER

Oral Suspension

Anthelmintic for Sheep, Goats, Cattle, Camels, Equines.



COMPOSITION: Each ml contains:

Fenbendazole

50 mg

PROPERTIES:

- An effective, safe and broad spectrum anthelmintic of the benzimidazole group.
- It has vermifugal, larvicidal and ovicidal activity against nematodes (Round Worms) and cestodes (Tape worms).

INDICATIONS:

For treatment and control of parasitic gastro – enteritis and verminous bronchitis.

■ Sheep, Goats, Cattle, Camels:

- **Gastrointestinal Round Worms:** Ostertagia, Trichostrongylus, Haemonchus, Strongyloides, Nematodirus, Cooperia, Bunostomum, Oesophagostomum, Marshallagia, Chabertia...
- **Lung Worms:** Dictyocaulus, Muellerius.
- **Tape Worms:** Moniezia.

■ Equines: Small and large Strongyles, Oxyuris, Ascarids, Trichostrongylus, Strongyloides.

DOSAGE: (single dose).

- **Shake well before use.**
- **Sheep, Goats:** 1 ml / 10 kg B. wt. (5 mg / kg B. wt.)
- **Cattle, Camels:** 1.5 – 2 ml / 10 kg B. wt. (7.5 mg / kg B. wt.)
For tape worms in cattle: 2 ml / 10 kg B. wt.
(10 mg activity / kg B. wt.)
- **Equines:** 1-2 ml / 10 kg B. wt. (5mg activity / kg B. wt.)
For ascarids in equines: 2 ml / 10 kg B. wt.
(10 mg / kg B. wt.)

SAFETY PERIOD:

- For meat consumption: 14 days.
- For Milk consumption: 3 days.

REMARKS:

- Repeat treatment after 4-6 weeks, when necessary.
- Regular deworming at intervals of 6-8 weeks may be required for equines.

STORAGE: Keep in a dry, dark place below 30 °C .

PACKING: (100, 250, 500, 1000) ml.