

DIAZINON-5%

INSECTICIDE
GRANULES

VAPCO

Class:
Organophosphate

Non-systemic insecticide with a wide spectrum uses in agricultural field with contact and stomach action.

COMPOSITION: Each kg contains:

DIAZINON

5% W/W (A.I)

PROPERTIES:

DIAZINON-5%: Non-systemic insecticide with contact, stomach and respiratory action on soil insects.

DIAZINON-5%: Moderately toxic to mammals.

DIAZINON-5%: Control a wide range of soil insect and worms which attacking agricultural crops, like Wireworms, Mole crickets, Rootworm, Beetles, Stem borer ... etc.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE KG / DONUM	NOTES
Potato, Tomato, Squash, Cucumber, Cabbage, Watermelon, Onion.	Cutworms	2.5	Scattering on soil
Apple, Pear, Plum, Apricot, Peaches, Nectarine	Tropinota Squalida Larvae	3	Scattering on soil
Olives	Borers, Soil Insects	2 - 3	Scattering on soil
Maize	Rootworm, Wireworm, Stem borer, Diptera Larvae.	4	Used Once by Scattering on soil
Cereals	Cutworms, Stem Borer, Beetles.	1 – 2	---
Rice	Leafminers	2.8	Scattering on soil
Soybean	Cutworms	2.4	Scattering on soil
Cotton	Cutworms	2.4	Scattering on soil
Peanuts	Cutworms	2.4	Scattering on soil
Sugarcane	Sugarcane beetles (<i>Pentodon Bispinosus</i>)	2.8	Scattering on soil
Gardens and Lawns	Beetles, Wireworm, Ants, Borers	10	Distributed evenly in the soil while applied through irrigation water to the Lawns.

Donum = 1000 m²

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

COMPATIBILITY:

Compatible with most pesticides of similar presentation but incompatible with alkaline compounds.

*** OTHER FORMULATION & TRADE NAME: DIAZINON - 60% EC, DIAZINON - 10% GR, VAZINON - 60% EC, VAZINON - 10% GR**

• For more details about first aid & precautions please refer to first aid & precautions index.