# **DINOMITE-20%**

## ACARICIDE/ INSECTICIDE WETTABLE POWDER



Highly effective to control all developing stages of mite, also controls Aphids, Whitefly, Thrips, and Leaf hoppers,

**COMPOSITION:** Each Kg contains:

PYRIDABEN 20% W/W (A.I)

PROPERTIES:

DINOMITE-20%: Highly effective to control all developing stages of mites (eggs, larvae, nymph

and adult).

DINOMITE-20%: Is effective to control insect pests such as Aphids, Thrips, Leaf hoppers and

Whitefly.

DINOMITE-20%: Rapid knockdown effect and long residual activity.

**DINOMITE-20%:** Temperature and pH do not affect the activity of the pesticide.

#### **USES AND APPLICATION RATE:**

CROP	PESTS	APPLICATION RATE GM / 20 LT. WATER	SAFETY PERIOD (DAYS)
Apple, Cherry	European red spider mite, Two-spotted spider mite, Apple rust mite, Red citrus spider mite, Citrus rust mite, Strawberry mite, Aphid, Whitefly, Thrips, Leaf hoppers.	15 - 20	21
Pear	European red spider mite, Two-spotted spider mite, Apple rust mite, Red citrus spider mite, Citrus rust mite, Strawberry mite, Aphid, Whitefly, Thrips, Leaf hoppers.	15 - 20	14
Apricot, Plum	European red spider mite	15	14
Peach	European red spider mite, Two-spotted spider mite, Apple rust mite, Red citrus spider mite, Citrus rust mite, Strawberry mite, Aphid, Whitefly, Thrips, Leaf hoppers.	15 - 20	14
Almond, Pomegranate	European red spider mite, rust mite, Aphid	15 - 20	21
Citrus	European red spider mite, Two-spotted spider mite, Apple rust mite, Red citrus spider mite, Citrus rust mite, Strawberry mite, Aphid, Whitefly, Thrips, Leaf hoppers.	5 - 10	3

<sup>•</sup> For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE GM / 20 LT. WATER	SAFETY PERIOD (DAYS)
Grapes	European red spider mite, Leaf hoppers.	15 - 20	45
Tomato, Cucumber, Eggplant	European red spider mite, Two-spotted spider mite, Whitefly.	15 - 20	7
Date Palm	European red spider mite, rust mite, Aphid.	15 - 20	21
Ornamental Plants	European red spider mite, Whitefly.	15 - 20	

#### CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

### **RE-ENTRY PERIOD:**

12 hours after the last application.

#### COMPATIBILITY:

Compatible with most pesticide except alkaline compound.

<sup>•</sup> For more details about first aid & precautions please refer to first aid & precautions index.