


DO – CARB

Anti-Protozoal
Injectable Solution



COMPOSITION: Each ml contains:

Imidocarb Dipropionate 120 mg

INDICATIONS:

- Do-carb is used for the control of Babesiosis in cattle, sheep, equines and dogs.
- It's also used for the treatment & prevention of Anaplasmosis in cattle and Ehrlichia canis infections in dogs.

DOSAGE:

Treatment:

- **Cattle:**(by subcutaneous injection)
 - For Babesiosis: 1 ml per 100 kg B.wt.
 - For Anaplasmosis or mixed infection due to Anaplasma & Babesia: 2.5 ml per 100 kg B. wt.
- **Sheep:** (by subcutaneous injection)
 - For Babesiosis: 0.5 ml per 50 kg B.wt
- **Equines:** (by intramuscular or subcutaneous injection)
 - For Babesiosis: 2 ml per 100 kg B.wt.
 - B. equi infection may require 2 doses at an interval of 24 hrs.
- **Dogs:** (by subcutaneous injection)
 - For Babesiosis: 0.25 ml per 10 kg B.wt.
 - For Ehrlichiosis or mixed infection due to Ehrlichia & Babesia: 0.5 ml per 10 Kg B. wt 2 doses at an interval of 14 days.

Prevention:

Single dose will prevent Babesiosis for up to 4 weeks.

- **Cattle:** 2.5 ml per 100 kg B.wt.
- **Equines:** 2 ml per 100 kg B.wt.
- **Dogs:** 0.5 ml per 10 Kg B.wt.

REMARKS:

- Don't administer intravenously.
- Don't combine with choline-sterase inhibitors or pesticides.
- Don't use in lactating animals.

WITHDRAWAL PERIOD:

- For meat: 28 days.

STORAGE: keep in dry, dark place below 30⁰ C.

PACKING : (50, 100) ml.