

# DOCLIN

Water Soluble Powder

**VAPCO**

ANTIBACTERIAL

**COMPOSITION: Each gm contains:**

<b>Doxycycline HCL</b>	<b>300 mg</b>
<b>Colistin Sulphate</b>	<b>800,000 I.U.</b>

**PROPERTIES:**

- Doxycycline is broad spectrum, long acting antibiotic, rapidly absorbed and slowly excreted.
- Colistin is very active against Gm-ve bacteria and specially the enteric infections.
- The combination of both antibiotics results in a strong antibacterial action which enhances the efficacy against the enteric pathogens as well.

**INDICATIONS:**

- For Poultry, game – birds, pigeons, calves, lambs and kids.
- Recommended for the different infections of the respiratory and alimentary tracts such as: CRD, Coryza, E. coli, Mycoplasmosis, Air sacculitis, Salmonella, Fowl cholera, Necrotic enteritis (Clostridia), Omphalitis and yolk sac infection, bacterial enteritis and diarrhea.
- Preventive therapy against early mortality of the newly – born and young animals.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

**DOSAGE:**

- **Treatment:** (Duration for: 3-5 days ).
  - **Poultry:** 100 gm / 200 litres of drinking water.
  - **Game – birds, Pigeons:** 0.5 gm per litre of drinking water.
  - **Calves, Lambs, Kids:** 1-2 gm / 10 kg B.wt.daily and orally with drinking water.
- **Prevention:** Half dosage for the same perios.

**SAFETY PERIOD:** (For meat consumption)

For calves, lambs, kids : 14 days.

For poultry : 7 days.

**REMARKS:**

- Not recommended for table-eggs layers.
- During treatment, mineral salts should not be added to the medicated water.
- During treatment, eggs should not be used for human consumption

**STORAGE:** Keep in a dry, dark place below 30 °C.

**PACKING:** (100, 250, 500, 1000) GM.