


# DOXYVAP

Water Soluble Powder



**COMPOSITION: Each gm contains:**  
**Doxycycline HCL                      200 mg**

**PROPERTIES:**

- Doxycycline, the broad spectrum antibiotic, is one of the tetracycline group; which is more rapidly absorbed in the gastrointestinal tract, distributed extensively throughout the body, and slowly excreted, than other tetracyclines.
- Doxycycline is a long acting antibiotic, and is a safe treatment even with impaired renal function. It is rapidly absorbed from the intestine and slowly excreted.
- Doxycycline is used in cases of different infections of respiratory, urogenital and gastrointestinal tracts; which are caused by mycoplasma and various gm+ve and gm-ve bacteria (Staph, Strept, Coli, Chlamydia, Clostridia, Rickettsia, Vibrio, Campylobacter, Pasteurella, Leptospira, Spirochaeta, Pseudomonas, Listeria, Shigella, Klesbsiella etc...)

**INDICATIONS:**

- Given to poultry, pigeons, caged-birds, calves, lambs, kids, dogs, cats and pigs.
- For the treatment and prevention of the different diseases: CRD, Coryza, Mycoplasmosis, Air succulitis, Sinusitis, Synovitis, Colibacillosis, Omphalitis and yolk sac infection, Fowl cholera, Necrotic enteritis (Clostridia), and different bacterial enteritis and diarrhea.

**DOSAGE:** (Treatment duration: 4-5 days)

- **Treatment:**  
Poultry: 100 gm / 200 litres of drinking water.  
Pigeons, Caged birds: 0.65 gm / litre of drinking water.  
Different animals and pigs: 0.5gm / 10 kg B.wt. daily and (orally with water).
- **Prevention:** Half dosage for the same period.

**SAFETY PERIOD:**

- For meat consumption : calves, lambs, kids : 14 days.  
Pigs : 8 days.  
Poultry : 5 days.
- For eggs consumption : 4 days.

**REMARKS:**

- During treatment, minerals salts should not be added to the medicated water.

**STORAGE:** Keep in dry, dark place below 30 °C.

**PACKING:** (100, 250, 500, 1000) GM.