

EL-TORO 25%

**INSECTICIDE
SUSPENSION CONCENTRATE**

VAPCO

Class:
Neonicotinoid

Highly effective Systemic Insecticide with contact and stomach action, to control many of sucking and chewing insects on a wide range of crops listed in the table.

COMPOSITION: Each liter contains:

THIAMETHOXAM

25% W/V (A.I)

PROPERTIES:

EL- TORO 25% : Highly effective systemic insecticide with contact and stomach action.

EL- TORO 25% : Highly effective with low dosage and long lasting effect.

EL- TORO 25% : Has a special formula that interferes with information transfer between nerve cells making the insects become paralyzed.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Cucumber, Muskmelon, Pumpkin, Squash, Watermelon	Aphids	2 – 4	7
	Cucumber Beetles, Leafminer, Whiteflies.	4 – 8	
Eggplant, Pepper, Tomato	Aphids, Colorado Potato Beetle, Flea Beetles, Leafhoppers.	3 – 4	7
	Whiteflies, Pepper Weevil, Stink Bugs.	4 – 8	
Lettuce, Celery, Parsley, Spinach	Aphids, Flea Beetles, Leafhoppers	2 – 4	7
	Whiteflies	4 – 8	
Cabbage, Cauliflower, Brussels Sprouts, Broccoli	Aphids, Flea Beetles	2 – 4	7
	Thrips, Whiteflies	4 – 8	
Mint	Aphids, Mint Flea Beetles, Leafhoppers	2 – 4	7
Carrot, Garden Beet, Turnip	Aphids, Flea Beetles, Leafhoppers	2 – 4	7
	Whiteflies	4 – 6	
Radish	Aphids, Flea Beetles, Leafhoppers	2 – 4	7
	Whiteflies	4 – 6	
Okra	Aphids, Whiteflies	2 – 4	5
Cassava, Potato, Sweet Potato, Yams	Colorado Potato Beetle, Flea Beetles, Potato Leafhoppers	2 – 4	14
	Aphids	4	

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Barley	Aphids	6	21
Rice	Stem Borer, Thrips Leaf folder, Brown Planthoppers, White Backed Hoppers, Green Leafhoppers,	2 – 4	14
Cotton	Aphids, Thrips	2 – 4	21
Lemon, Orange, Pummelo, Mandarin, Grapefruit	Aphids, Leafhoppers	4 – 6	14
	Asian Citrus Psyllid, Citrus Black Fly, Citrus Leafminer, Mealybugs, Soft Scales, Whiteflies	6 – 8	
	Citrus Root Weevil Adult, Fruit Fly, Grasshoppers, Plant Bugs, Stink Bugs, Thrips (foliage feeding)	7 – 8	
Pome Fruit: Apples, Loquat, Quince	(Pre-bloom): Apple Aphid, Leafminers, Bugs.	7	35
	(post-bloom): Leafhoppers	3 – 4	14
	(post-bloom): Aphid, Leafminers, Plum Curculio	7 – 8	35
Pome Fruit: Pears	(Pre-bloom): Pear Psylla, Apple Aphid	7 – 8	35
	(post-bloom): Apple Aphid, Mealybug, Plum Curculio, Pear Psylla	7 – 8	35
	(post-bloom): Leafhoppers	3 – 4	14
Mango, Avocado, Papaya, Sapodilla	Aphids, Leafhoppers, Mealybugs, Whiteflies, Thrips (foliage feeding)	6	---
Apricot, Peach, Nectarine, Plum	Leafhoppers	3 – 4	14
	Aphids	4 – 6	
	Cherry Fruit Fly, Stink Bugs, Thrips.	7 – 8	
Grape	Japanese Beetle, Leafhoppers, Mealybugs	2 – 5	5
Strawberry	Aphids, Leafhoppers	2 – 4	3
	Bug, Adult Weevil	6	
Tea	Mosquito Bug	2 – 4	7

CAUTION:

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- Do not spray during flowering stage.

COMPATIBILITY:

Compatible with most pesticide except alkaline and oxidative materials.

RE-ENTRY PERIOD:

24 hours after the last application.

- For more details about first aid & precautions please refer to first aid & precautions index.