EMAVAP-2%

INSECTICIDE EMULSIFIABLE CONCENTRATE



Non-systemic insecticide which penetrates leaf tissues by translaminar movement, Shortly after exposure to EMAVAP-2% affected larvae are paralyzed, stop feeding and subsequently die after 2-4 days.

COMPOSITION: Each liter contains:

EMAMECTIN BENZOATE

2% W/V (A.I)

PROPERTIES:

EMAVAP-2%: Has contact activity, but is most efficacious when ingested by the pest.

EMAVAP-2%: Non-systemic insecticide for the control of the larval stages of certain Lepidoptera

attacking a wide spectrum crops, it is active against insects which have become

resistant to other traditional insecticides.

EMAVAP-2%: Penetrates leaf tissues, which mean a long lasting residual effect with guick killing.

EMAVAP-2%: Non-phytotoxic to plants.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	COMMENTS	SAFETY PERIOD (DAYS)
Cabbage, Broccoli, Cauliflower	Beet armyworm, Diamondback moth, cabbage moth.	4.25 - 8.5		14
	Cabbage looper, Leafminer	5.6 - 8.5		7
Lettuce, Celery,	Beet armyworm, Tobacco budworm.	4.25 - 8.5	Apply when larvae are first observed. Additional applications may be repeated to maintain control.	7
Spinach	Cabbage looper, Leafminer	5.6 - 8.5		
Eggplant, Pepper, Tomato	Beet armyworm, Cabbage looper, Tobacco budworm, Tomato pinworm. Alfalfa looper, Leafminer Tuta Absoluta	4.25 - 8.5 5.6 - 8.5 10		7
Watermelon	Thrips	12.5 - 15		
Citrus	leafminers	12.5 - 15		
Apple, Pear, Quince	Bud moths, Leafminers, Leafrollers, Spider mite	4.25 - 8.5		14
Maize	Maize worm	1		7
Rice	leafminer	10 - 12		
Tobacco	Bud moths, Podworm	1		7

[•] For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

72 hours after the last application.

COMPATIBILITY:

Is compatible with most insecticides, fungicides, and foliar nutrient products, but it is favorable to make compatibility test before use.

^{*} OTHER FORMULATION: EMAVAP - 5% SG.

[•] For more details about first aid & precautions please refer to first aid & precautions index.