ERYTHRODONE

Injectable Solution For Sheep, Goats, Cattle, Pigs



COMPOSITION: Each ml contains:

Erythromycin Thiocyanate 100 mg Lidocaine Hydrochloride 50 mg

PROPERTIES:

- Erythromycin, a macrolide antibiotic which has a broad spectrum activity, outstanding therapeutic
 effectiveness and safety.
- It is highly active against gm+ve and gm-ve bacteria: Staph, Sterpt, Pneumococci, Pasteurella, Haemophilus, Brucella, Mycoplasma, Listeria, Rickettsia, Clostrdia, Campylobacter, Entorococci, Erysiplothrix.
- It reduces the virulence of bacteria, even those that are not included in its spectrum of activity.

INDICATIONS:

- Respiratory tract infections:
 - Tracheitis, Bronchitis, Bronchopneumonia, Pneumonia, Rhinitis,
- Gastrointestinal tract infections:
 - Entertis, Lamb dysentery (Clostridia), peritonitis, Reticulo-peritonitis.
- Urogenital tract infections:
 - Nephritis, Pyelonephritis, Cystitis, Metritis, Vaginitis.
- Mastitis (supportive treatment), Foot Rot, Mycoplasmosis, Shipping fever, Septicaemia, Leptospirosis, Pyeamia, Erysipelas, Vibriosis.
- Skin infections: Pyoderma, Furunculosis, Abscesses.

DOSAGE: Deep intramuscular injection:

■ Cattle, Sheep, Goats, Swine: 1 ml / 20 kg B.wt. given once or twice daily for 2-3 days.

REMARKS:

- Change the injection site each time, and the dose may be subdivided.
- A transient local irritation may occur after injection for few days.
- Not given with penicillins, as interference action may occur.

SAFETY PERIOD:

For milk consumption: 3 days. For meat consumption: 7 days.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (50, 100) ml.