

FAST-5%

INSECTICIDE

EMULSIFIABLE CONCENTRATE

VAPCO

Class:
Pyrethroid

A broad spectrum pyrethroidal insecticide based on ESFENVALERATE as active ingredient.

COMPOSITION: Each liter contains:

ESFENVALERATE

5% W/V (A.I)

PROPERTIES:

FAST-5%: Contains ESFENVALERATE which has a wide spectrum activity on agricultural insects.

FAST-5%: Has special activity against insect that gain resistant to Organophosphorus, Organochlorine and Carbamate compounds.

FAST-5%: Effective against Coleoptera, Hemiptera, Diptera, Lepidoptera and Orthoptera.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Tomatoes	Fruitworm	3 - 8	1
Cabbage	Cutworm		3
Cucumber	Aphid		3
Muskmelon	Whitefly		3
Potatoes, Eggplant	Thrips, Green Peach Aphid, Colorado Potato Beetle	5 - 8	7
Carrots	Cutworm, Leafhopper	22 - 37	7
Broccoli	Alfalfa looper, Cabbage looper, Cutworm, Flea beetle, Grasshopper	22 - 37	3
Apples	Psylla	6 - 8	21
	Wooly aphids	4 - 5	
Peach, Almonds, Apricots	Fruit fly, Piercing pests	4 - 6	21
Sugar Beet	Cabbage Moth, Cutworm, Aphids	4 - 6	21
	Mangold Flea Beetle, Beet Tortoise Beetle	4 - 6	
Sugarcane	Sugarcane borer	22 - 37	21
Peas, Beans	Pea Moth, Pea Aphid, Black Bean Aphid, Weevil, Pea Thrips	4 - 6	3
	Pea Leaf Weevil	3 - 4	

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Cereals	Worm	6 – 8	1
	Aphid	2 - 4	
	Sunn Pest	4 - 6	
	Cereal Leaf Beetle	2 - 5	
Maize	Corn Earworm	4 - 6	---
	Corn Borer, Fruit Fly	6 - 8	
Sorghum	Aphids	5	---
Soybean	Green cloverworm, Mexican Bean beetle, Potato leafhopper, Saltmarsh caterpillar, Valvetbean caterpillar, Woollybear caterpillar	22 - 37	21
	Bean Leaf Beetle, Cabbage Looper, Corn Earworm, Cutworms, Grasshoppers, South Green stink bug, Three-cornered alfalfa hopper	22 - 37	
Peanuts	Corn earworm, Potato leafhopper, Red-necked peanut worm, Valvetbean caterpillar	11 - 22	21
	Cutworm, Granulate cutworm, Grasshopper	22 - 37	
	Black cutworm, Chinch bugs, Cutworms	22 - 37	
Pecans	Aphids, Nut casebearer leaf phylloxera, Stem phylloxera, Weevil	19 - 57	21
Walnuts	Codling moth, Navel orangeworm, Aphid, Husk fly	37 - 75	21
Artichoke	Plume moth	22 - 37	1
Cotton	American Bollworm, Cotton Bollworm, Tobacco Budworm, Spiny Bollworm, Spotted Bollworm, Common Cutworm, Cotton Leafworm	7.5 - 12	21
	Aphid	6 - 10	
	Whitefly, Leafhopper, Cotton Jassid, Cotton Boll Weevil, Thrips	10 - 12	
Oil Seed Rape	Cabbage Stem Weevil, Rape Stem Weevil, Turnip Gall Weevil, Cabbage Seed Weevil, Turnip Flea Beetle, Small Striped Flea Beetle, Large Striped Flea Beetle, Brassica Pod Midge	4 - 6	---
	Turnip Sawfly	2 - 4	

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

72 hours after the last treatment.

COMPATIBILITY:

Compatible with most pesticides except alkaline compounds.

*** OTHER FORNULATION: FAST – 7.5% ULV.**

• For more details about first aid & precautions please refer to first aid & precautions index.