

# FENATODE-10%

NEMATICIDE  
GRANULES

VAPCO

Class:  
Organophosphate

**Systemic nematicide with contact action used to control Nematodes in Tomato, Citrus, Banana, and Potato field.**

**COMPOSITION:** Each kg contains:

**FENAMIPHOS**

**10% W/W (A.I)**

**PROPERTIES:**

**FENATODE-10%:** Highly effective Organophosphorus nematicide with cholinesterase inhibitor activity.

**FENATODE-10%:** Systemic nematicide with contact action, absorbed by the roots, and translocated to the leaves.

**FENATODE-10%:** Readily distributed in the soil so that it controls Nematodes present in deeper soil layers. Also It has a residual activity for several months.

**FENATODE-10%:** Stimulates growth, increase yield and improve quality.

**FENATODE-10%:** Effective against soil pests like populations of ectoparasitic, endoparasitic, Free-living, Cyst-forming and Root-knot nematodes.

**USES AND APPLICATION RATE:**

CROP	PEST	DOSAGE	SAFETY PERIOD (DAYS)
<b>Citrus</b> * Small trees * Old trees	Slow Decline Nematode	Used once in spring in rate 40 gm / m <sup>2</sup> . Add 20 - 40 gm / m <sup>2</sup> after harvest immediately, in case of sever infestation apply again with the same rate in autumn by applying under trees and mix to depth 5 - 10 cm and irrigate in rate 25 lt. / m <sup>2</sup> .	90
<b>Tomatoes</b>	Nematode	In open filed apply on rows before 14 days before seedling plant in the soil, howing to depth 5 - 10 cm then irrigate heavily 25 lt. water/m <sup>2</sup> in rate of 1.5 - 3 kg./donum. In green house apply 14 days before planting in the rate of 4.5 kg/donum, howing to depth 5-10 cm and irrigate heavily.	90
<b>Potatoes</b>	Nematode	Apply in the row with tubers then cover the row and irrigate heavily in a rate of 1.5 kg/donum. To treat the whole field apply in rate 4 - 6 kg./donum two weeks before planting tubers.	60
<b>Banana</b>	Nematode	New shoots: cover shoots with mixture of clay and water in addition to <b>FENATODE-10%</b> in the rate of 8 kg./100 lt. Clay, retreat after 4 months in sever infection by application of granules 25 - 30 cm away from the hole during boot stage then irrigate heavily.	60

Donum = 1000 m<sup>2</sup>

• For more details about first aid & precautions please refer to first aid & precautions index.

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest

**COMPATIBILITY:**

Compatible with the same formulation, incompatible with alkaline materials.

• For more details about first aid & precautions please refer to first aid & precautions index.