

FLUZURON-15%

**INSECTICIDE
INSECT GROWTH REGULATOR (IGR)
SUSPENSION CONCENTRATE**

VAPCO

**Class:
Benzoylurea**

Systemic insect growth regulator begins its effectiveness at the time of moulting or oviposition

COMPOSITION: Each liter contains:

TEFLUBENZURON

15% W/V (A.I)

PROPERTIES:

FLUZURON-15%: Chitin synthesis inhibitor that interferes with the formation of insect cuticle and lead to disturbance in moulting process of the insect.

FLUZURON-15% : Economic pesticide effective at a lower dose.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Potato, Tomato, Eggplant, Pepper, Cucumber	Leafhopper, Beetles, Lepidoptera, Cutworm, Crawling Worm	10	3 days except potatos 28 days
Cabbage	Cabbage Butterfly, Cabbage Moth	3 - 4	7
Apple, Pears	Borers Fruit Moth Psylla	6 - 8 6 - 10 10	14
Plums, Peaches, Nectarines, Apricot, Cherries, Almond	Fruit Moth Borers	6 - 10 6 - 8	21
Sugar Beet	Cotton Leafworm	10	28
Cotton	Cutworm	7 - 12	28
Ornamental Plants	Autumn Moth	10	---
Pasture	Leafhopper, Beetles, Lepidoptera, Cutworm, Crawling Worm	10	3

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- In case of severe infection repeat after 7 days from the first spray.

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY:

Compatible with most pesticides except highly acidic and oxidizing compounds, but it is preferable to make mixing trial before use.

*** OTHER FORMULATION: FLUZURON-5% ULV.**

• For more details about first aid & precautions please refer to first aid & precautions index.