

# FOLDAN-50%

**FUNGICIDE  
WETTABLE POWDER**

**VAPCO**

Class:  
Phthalimide

**A protective fungicide with contact action, effective in controlling many of the most fungal diseases on a wide range of crops.**

**COMPOSITION:** Each kg contains:

**FOLPET**

**50% W/W (A.I)**

**PROPERTIES:**

**FOLDAN-50%:** A contact fungicide with a curative and protective action.

**FOLDAN-50%:** Effective to control many of the most economically devastating fungal diseases such as: Downy Mildew, Powdery Mildew, Leaf Spot, Scab,.... etc.

**FOLDAN-50%:** Used on a wide range of crops such as Pome fruit, Stone fruit, Citrus. Grapes, Olives, Potatoes, Cabbage, Onions, Tomatoes, Cucumber, Eggplant,.... etc.

**FOLDAN-50%:** Slightly toxic to mammals and other beneficial insect or wildlife.

**USES AND APPLICATION RATE:**

CROP	DISEASE	APPLICATION RATE GM / 20 LT. WATER	NUMBER OF APPLICATION	SAFETY PERIOD (DAYS)
Pear	Scab, Leaf Spot, Powdery Mildew	50	10	21
Apple		40	Every 14 days	21
Citrus	Anthraxnose, Dieback, Gummosis	60	---	21
Grapes	Downy Mildew, Powdery Mildew, Brown Rot, Dieback.	60	Every 14 days	8
Apricot, Plum, Peach, Almond	Leaf Curl, Anthracnose, Powdery Mildew, Leaf Spot.	50	3	10
Bush fruit (Blueberries, Current, Gooseberries) Cane fruit (Blackberries, Loganberries, Raspberries, Youngberries)	Botrytis, Leaf Spots	30 - 40	Spray when flowering. Repeat at 7 – 14 day intervals from flowering until harvest. Use the higher rate when conditions favour infection.	---

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE GM / 20 LT. WATER	NUMBER OF APPLICATION	SAFETY PERIOD (DAYS)
<b>Olives</b>	Peacock Eye Spot, Wilt	30	---	10
<b>Cucurbits</b>	Powdery Mildew, Downy Mildew, Stem Rot, Anthracnose.	15	---	7
<b>Eggplant, Peas</b>	Powdery Mildew, Early Blight, Late Blight	30	---	10
<b>Legumes</b>	Powdery Mildew, Leaf Spot	7 - 8	---	10
<b>Lettuce</b>	Downy Mildew, Gray Mold	40	Weekly	11
<b>Muskmelon</b>	Downy Mildew, Powdery Mildew, Stem Rot, Anthracnose.	40	Weekly	11
<b>Watermelon</b>	Downy Mildew, Powdery Mildew, Stem Rot, Anthracnose.	50	1 - 2	7
<b>Onions</b>	Pink Blotch, Neck And Leaf Rot.	40 - 50	2 - 3	7
<b>Pepper</b>	Wilt, Stem Rot, Powdery Mildew.	10 - 30	3 - 4	7
<b>Potatoes</b>	Early Blight, Late Blight	60	Weekly	14
<b>Tomatoes</b>	Early Blight, Late Blight	40	Weekly	14
<b>Roses</b>	Black Spot, Powdery Mildew	40	Apply at regular intervals from budburst.	---

#### CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

#### PHYTOTOXICITY:

Not phytotoxic, except for: Sweet Cherries, D'Anjou variety of Pears and Russetting is possible in sensitive Apple varieties if applied early in cropping.

#### RE-ENTRY PERIOD:

24 hours after the last application.

#### COMPATIBILITY:

Compatible with most pesticides except highly alkaline compounds. It's preferable to use the product immediately after mixing.

• For more details about first aid & precautions please refer to first aid & precautions index.