HERBIFLURIN-48%





A selective pre-planting soil-herbicide to control Annual grasses and Board-leaved weeds and used to control weed seeds as they germinate

COMPOSITION: Each Liter contains:

TRIFLURALIN 48% W/V (A.I)

PROPERTIES:

HERBIFLURIN-48%: A selective pre-planting soil-herbicide.

HERBIFLURIN-48%: Has a good efficiency to control annual narrow and broad-leaved weeds.

HERBIFLURIN-48%: Has low toxicity to mammals.

HERBIFLURIN-48%: Has no residual risk on the crop leaves, fruits or seeds, also it has no

injury on the second planting crop.

USES AND APPLICATION RATE:

Must be applied and incorporated before planting and emergence and well mixed with soil, Treated soil should be wet or irrigate after the application.

CROP	METHOD OF APPLICATION	RATE OF APPLICATION	APPLICATION TIME
A pre-planting herbicide for the following crops:	Apply before planting by mixing with the	Applied at the Rate of 120-250 ml	
Seedling planting: Pepper, Tomatoes, Tobacco, Cauliflower, Cabbage, Broccoli.	soil at a rate of 20-40 liter of spraying solution / donum to the depth 8-15 cm.	/ donum / 20 - 40 lt water.	or transplanting, also it can be applied 90 days after planting but it must avoid crops contact.
Seed planting: Soybean, okra, carrots, turnip.			
Sugar beet (seedling) (spray between planting lines after the emergence of 4-6 leaves).			

CAUTION: Donum = 1000 m²

- Used only on above mentioned crops.
- Do not plant the following crops before 6-12 months, after spraying: Muskmelon, Watermelon, Cucumber, Spinach, Corn, Tomato (seeds), Pepper (seeds), Cauliflower (seeds), Cabbage (seeds) and Sugar beet (seeds).

GRAZING RESTRICTIONS: Do not graze or feed animal from treated area for 7 days at least.

RE-ENTRY PERIOD: 24 hours after the last application.

COMPATIBILITY: It is preferable not to be mix with other pesticides.

• For more details about first aid & precautions please refer to first aid & precautions index.