

# HERBSTAR-25%

HERBICIDE  
EMULSIFIABLE CONCENTRATE

VAPCO

Class:  
Oxadiazole

A selective, pre-emergence herbicide for controlling many annual grasses and broad-leaved weeds.

**COMPOSITION:** Each Liter contains:

**OXADIAZON**

**25% W/V (A.I)**

**PROPERTIES:**

**HERBSTAR-25%:** A selective, pre-emergence herbicide to control many annual grasses and broadleaf weeds.

**HERBSTAR-25%:** Is absorbed mainly through the leaves and is translocated rapidly, inhibiting cell division in root and shoot tips and stop growth within hours of treatment.

**HERBSTAR-25%:** One application provides season-long control of weeds.

**HERBSTAR-25%:** Dose not thermal and photo-decompose or volatilize and will remain on the soil surface for long periods.

**USES AND APPLICATION RATE :**

A selective contact pre-emergence herbicide, absorbed by foliage and is rapidly move to the growing tips of roots and foliage soon after application (foliage system, root system).

CROP	WEEDS	TIME OF APPLICATION	APPLICATION RATE
Onions	Broad-leaved weeds such as: Crown daisy, Viscum, Nettle, Wild mustard, Naked launaea, Galium and	Spray early when the plants 15 cm tall (after 3 leaf stage) or after 20-30 days from transplanting, or spray immediately after seeding.	200 - 300 ml / donum depends on weeds density.
Garlic	Annual grasses such as : Shepherd's purse, Aleppo Spurge, Wild oat, Bristly foxtail	Spray after planting the bulbs before the weeds germinate it is advisable to irrigate in order to increase the activity of the herbicide.	
Broadbeans, Grapes, Apricot, Peach, Plum, Cherry, Nectarine, Apple, Pear and Ornamentals		Pre-emergence.	

Donum = 1000 m<sup>2</sup>

**CAUTION:** Used only on above mentioned crops and follow the recommended re-entry period and grazing restrictions.

• For more details about first aid & precautions please refer to first aid & precautions index.

**REMARKS:**

- Calibration of sprayer is important before application.
- Avoid hoeing the soil immediately after treatment.
- The soil must be wetted at the time of spraying.
- **HERBSTAR-25%** don't control weeds that propagate by bulb's or rhizomes.
- Results of control appears after 5 - 10 days from spraying.
- No residues will be left on Onions during harvest.

**GRAZING RESTRICTIONS:**

Do not graze or feed animal from treated area.

**RE-ENTRY PERIOD:**

24 hours after the last application.

**\* OTHER TRADE NAME: ROOTAGE-25% EC.**

• For more details about first aid & precautions please refer to first aid & precautions index.