

# JAZZAR-25%

ACARICIDE  
WETTABLE POWDER

VAPCO

Class:  
Organotin Acaricide

**Broad spectrum acaricide against all motile stages of mites.**

**COMPOSITION:** Each Kg contains:

**AZOCYCLOTIN**

**25% W/W (A.I)**

**PROPERTIES:**

**JAZZAR-25%:** long-acting acaricide properties, stable to light and rain.

**JAZZAR-25%:** long- acting acaricide with contact action to control all motile stages of mites on fruit trees, vegetables and ornamentals.

**USES AND APPLICATION RATE:**

CROP	PESTS	APPLICATION RATE GM / 20 LT. WATER	REMARKS	SAFETY PERIOD (DAYS)
Beans, Okra, Peas	European Red Mite, Two-Spotted Spider Mite.	20 - 25	---	7
Tomatoes, Eggplant, Potatoes, Cucumbers, Squash, Peppers, Cabbage, Cauliflower, Muskmelon, Watermelon, Onions, Garlic	European Red Mite, Two-Spotted Spider Mite.	20 - 25	---	7
Apple, Pear	European Red Mite, Two-Spotted Spider Mite.	10 - 15	A) Early Season Apply as a dilute spray to coincide with the mite hatchings	15
		15 - 20	B) Mid and Late Season Apply as a dilute spray to coincide with mite hatchings	
Plums, Peaches, Nectarines, Apricot, Cherries, Almond	European Red Mite, Two-Spotted Spider Mite.	10 - 15	Apply as a dilute spray to coincide with summer mite hatchings	15

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE GM / 20 LT. WATER	REMARKS	SAFETY PERIOD (DAYS)
<b>Currants Raspberries, Strawberry</b>	Two-Spotted Spider Mite.	15	Apply on first appearance of mites. A further application may be necessary 10-14 days later if mite numbers are high. Ensure complete coverage. Apply up to petal fall or post harvest.	7
<b>Citrus ( Lemon, Mandarin, Orange, Pummelo, Grapefruit)</b>	Citrus Red Mite	15	Apply at first appearance of mites. A further application may be necessary 10- 14 days later if mite numbers are high. Ensure complete coverage. Do not apply with oil or within 28 days of an oil application.	15
<b>Grapes</b>	Two-Spotted Spider Mite.	15 - 20	Apply at first appearance of mites. A further application may be necessary 10- 14 days later if mite numbers are high. Ensure complete coverage.	15
<b>Ornamental plant</b>	European Red Mite, Two-Spotted Spider Mite	20 - 25	---	---

**CAUTION:**

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- Don't use on roses and chrysanthemum

**RE-ENTRY PERIOD:**

48 hours after the last application.

**COMPATIBILITY:**

Compatible with most pesticide, except highly alkaline compound, but it is preferable to make mixing trial before use.

• For more details about first aid & precautions please refer to first aid & precautions index.