

# LIDOPHRINE

Injectable Solution  
Local anaesthetic

**VAPCO**  
INJECTABLE  
PRODUCTS

**COMPOSITION:** Each ml contains:

**Lidocaine hydrochloride**      **20 mg**  
**Epinephrine acid tartrate**      **0.01 mg**

**PROPERTIES:**

\* LIDOPHRINE is a fast acting local anesthetic that produces approximately twice the area of anesthesia produced by equal concentrations of procaine. epinephrine is added to prolong the duration of anaesthesia and to reduce systemic absorption.

**INDICATIONS :**

A local anaesthetic used in infiltration, nerve block and epidural anaesthesia in dogs, cats, cattle, horses and sheep.

**DOSAGE :**

<b>Horse:</b>	Volar nerve block	5 - 10 mL
	Low epidural (erect animal)	5 - 12 mL
	Infiltration	2 - 50 mL
<b>Cattle:</b>	Low epidural (erect animal)	5 mL
	N. Cornualis block (dehorning)	20 mL
	Paravertebral (at each site)	10 - 15 mL
	(for last thoracic)	20 mL
	Infiltration	5 - 100 mL
<b>Calf:</b>	Epidural	3 - 10 mL
<b>Sheep:</b>	Epidural	3 - 7 mL
	Infiltration	2 - 50 mL
<b>Dogs:</b>	Epidural	2 - 10 mL
	Brachial Plexus Block (average sized dog)	10 mL
	Infiltration	2 - 10 mL
<b>Cats:</b>	Epidural	0.5 - 2 mL
	Infiltration	0.5 - 4 mL

**NOTE:** The lowest possible concentration and dose should always be used. Inject slowly. Do not inject intravenously. Infiltration anaesthesia may be obtained by intramuscular or subcutaneous injection.

**REMARK:**

- Injections should be made immediately after withdrawal from the bottle, since the product may react with the metallic components of the syringe.



**VAPCO**  
INJECTABLE  
PRODUCTS

**WITHDRAWAL PERIOD:**

- Meat: 5 days before slaughtering
- Milk: 4 days.

**STORAGE:** keep in a dry, dark place below 30 °C .

**PACKING:** 100, 50 ml.