

MANTOX-FORTE

**FUNGICIDE
WETTABLE POWDER**

VAPCO

Class:
Dithiocarbamate ⁽¹⁾
Inorganic compound ⁽²⁾

Protective fungicide, highly effective in controlling many fungal diseases such as Blights, Downy mildew and many others on a wide range of agricultural crops.

COMPOSITION: Each kg contains:

MANCOZEB ⁽¹⁾	20% W/W (A.I)
COPPER IN THE FORM OF: (COPPER OXYCHLORIDE AND COPPER SULFATE) ⁽²⁾	21.5% W/W (A.I)
STIMULATING AGENT (IRON SULFATE)	6%

PROPERTIES:

MANTOX-FORTE: Protective fungicide, highly effective in controlling many fungal diseases such as Blights, Downy mildew and many others on a wide range of agricultural crops.

MANTOX-FORTE: With a broad spectrum action to control fungal diseases and protect from resistant formulation towards other pesticides.

MANTOX-FORTE: Has a longer protection period than any other fungicides, contains Mancozeb as active ingredient which is needed to the crop growth.

MANTOX-FORTE: Contains Zinc and Manganese which are needed to crop growth.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE GM/ 20 LT. WATER	SAFETY PERIOD (DAYS)
Cucumber, Tomato, Onion, Beans, Cabbage, Potato, Okra, Eggplant, Pepper	Blights, Anthracnose, Leaf Spots (<i>Cercospora</i>), Downy Mildew, Rusts, Angular Leaf Spot.	30 - 45	7
peas	Gray Mold, Leaf Spot, Downy Mildew, Anthracnose	40 - 60	7
Apple trees	Gray Mold, Leaf Spot (<i>Cercospora</i>), Downy Mildew, Anthracnose, Fruit Spot	40 - 60	7
Banana, Lemon, Orange, Pummelo, Mandarin, Mango, Almond, Peach, Apricot, Cherry, Nectarine	Leaf Spots, Fruit Spots, Anthracnose, Peach Leaf Curl.	30 – 45	21
Almond trees	Gray Mold, Leaf Spot (<i>Cercospora</i>), Downy Mildew, Anthracnose, Scab	40 - 60	7

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE GM/ 20 LT. WATER	SAFETY PERIOD (DAYS)
Guava	Gray Mold, Leaf Spot, Downy Mildew, Anthracnose, Scab	40 - 60	7
Cocoa	Gray Mold, Leaf Spot, Downy Mildew, Anthracnose.	40 - 60	7
Strawberries	Gray Mold, Leaf Spot, Downy Mildew, Anthracnose.	40 - 60	7
coffee	Gray Mold, Leaf Spot, Downy Mildew, Anthracnose.	40 - 60	7

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

TIME BETWEEN SPRAYS:

Repeated as needed.

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY:

Compatible with most pesticides, except Thiram, Lime sulfur, and Dithiocarbamates.

• For more details about first aid & precautions please refer to first aid & precautions index.