

MYCOPLASMIN

**Water Dispersible Powder
For Poultry & Other Animals**

VAPCO

ANTIBACTERIAL

COMPOSITION: Each gm contains:

Erythromycin Thiocyanate	250 mg
Trimethoprim	40 mg
Colistin Sulphate	500,000 I.U.

PROPERTIES:

- Erythromycin, a macrolide antibiotic, is safe and highly active against Mycoplasma, Gm+ve and Gm-ve bacteria such as:
Staph, Strept, Coryne, Coli, Pasteurella, Pneumococci, Haemophilus gallinarum, Erysipelothrix etc...
- Colistin is very active antibiotic against Gm-ve bacteria; and due to its slow and poor absorption from intestine, it concentrates its antibacterial action locally, and thus shows an outstanding effect against the enteric infections.
- Trimethoprim is an effective antibacterial agent, and has a reasonable antibacterial spectrum.

INDICATIONS:

- **Poultry:**
 - Cases of CRD, Mycoplasmosis, Coryza, Sinusitis, Air sacculitis, Synovitis, Salmonellosis, Pasteurellosis, Vibronic hepatitis, Colibacillosis, Clostridia, Staph and strept infections, Enteritis, Diarrhea etc...
 - A supportive treatment in cases of respiratory viral disease and clostridial infections.
 - Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.
- **Other animals:** Septicaemia and different infections of respiratory, alimentary and urogenital tracts: Pneumonia, Bronchitis, Pharyngitis, Enteritis, Bloody scour, Diarrhea, Nephritis, Cystitis etc...

DOSAGE:

Mix well before use.

- **Treatment: (use hot water 50-60 °C).**
 - **Poultry:** 100 gm per 200 litres of drinking water for 3-5 days .
 - **Other animals:** 1-5 gm / 25 kg B.wt. daily and orally with water for 3-5 days.
- **Prevention :** Half dosage for the same period.

SAFETY PERIOD: For meat and eggs consumption: 7 days.

STORAGE: Keep in dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.