


NEOCYCLIN

Water Soluble Powder



COMPOSITION: Each gm contains:

Oxytetracycline HCL	55.55 mg	Vitamin K3	0.5 mg
Neomycin Sulphate	55.55 mg	Vitamin B12	0.9 mcg
Vitamin A	1,120 I.U	Vitamin B2	0.002 mg
Vitamin D3	250 I.U	Nicotinic Acid	8.9 mg
Vitamin E	0.345 I.U	Ca- D - Pantothenate	3.4 mg

Trace elements salt : (Iron, Manganese, Zinc, Copper, Cobalt) 1.1 mg

PROPERTIES :

- Neocyclin represents a full treatment, thanks to a synergistic combination of antibacterials adapted to the therapy of the digestive tract.
- Broad spectrum activity of Oxytetracycline against various gram positive and gram negative bacteria, Spirochetes, Rickettsiae & Chlamydia.
- Bactericidal activity of neomycin in low concentration against certain gram positive and gram negative enteric pathogens including the important *Escherichia coli* organism.
- Antimicrobial activity of Oxytetracycline occurs both in the blood serum and gastro-intestinal tract, while the activity of Neomycin concentrates in the gastro-intestinal tract.
- The vitamins and trace elements contents activate, increase viability and support the body resistance against infections, to maintain optimum health.

INDICATIONS:

• Poultry

Infectious Enteritis , Fowl Cholera, Pullorum Disease, Fowl Typhoid, Monocytosis, Omphalitis, Hexamitiasis.

• Mammals

Infectious enteritis such as colibacillosis or salmonellosis, enterotoxaemia, secondary infections of viral diseases, bacterial respiratory diseases.

DOSAGE: (Duration for 3-5 days).

• Poultry:


- Treatment dose: 180 gm per 100 litres of drinking water.
- Prevention dose: Half dosage for the same period.

• mammals:

- Lambs, Kids: 2 gm per lamb daily with drinking water or milk.
- Calves: 14 gm per calf daily with drinking water or milk for the first 7 days of life.
- Beef cattle: 14 gm per beef cattle daily with drinking water for the first month of fattening period. At the last month of the fattening period repeat the same regimen.

SAFETY PERIOD:

For meat & eggs consumption: 7 days.



REMARKS:

- During treatment, mineral salts should not be added to the medicated water.

STORAGE: Keep in a dry, dark place below 25 °C.

PACKING: 180 GM.