

# OXYDONE-10%

Injectable Solution  
For livestock

**VAPCO**  
ANTIBACTERIAL/  
ANTIBIOTIC

**COMPOSITION:** Each ml contains:  
**Oxytetracycline - HCL** **100 mg**

## PROPERTIES:

- A broad spectrum antibiotic against susceptible Gm+ve bacteria (Staph, Strept, Coryne, Clostridia, Erysipelothrix) and Gm-ve bacteria (E.coli, Pasteurella, Actinobacillus, Moraxella), Mycoplasma, Protozoa, Haemophilus, Spirochaetes, Chlamydia, Leptospira, Rickettsia, Listeria and some large viruses.
- Distributed rapidly and extensively throughout the body tissues.

## INDICATIONS:

- For **Sheep, Goats, Cattle, Buffaloes, Camels, Pigs**.
- Treatment of different cases:
  - Shipping fever, Wooden Tongue (Actinobacillus), Pink eye (Bovine and ovine kerato conjunctivitis "Moraxella"), Swine Dysentery (Spirochaetosis), Atrophic Rhinitis in pigs, Dysagalactia, Foot rot, Calf diphtheria.
  - Anaplasmosis (Theileriosis) and Heart water in cattle (Ehrlichia "Rickettsia"), Red water of calves (Leptospira).
  - Enzootic abortion (Chlamydiosis) in sheep and goats.
  - Clostridial infections: Enterotoxaemia (Lamb dysentery, Black leg, Malignant oedema).
  - Scours, enteritis, diarrhea.
  - Septicaemia, Pyaemia, Peritonitis, Navel / Joint-III, wounds and skin infections, abscesses.
  - Supportive therapy in mastitis.
  - Respiratory tract infections: Bronchitis, pneumonia, rhinitis, etc.....
  - Urinary tract infections: Nephritis, Pyelonephritis, Cystitis.
  - Genital tract infections: Metritis, Puerperal Sepsis, etc...
  - Control of post-operative and post-parturient infections.
  - Bacterial complications of viral diseases.
- **DOSAGE:** (By deep intramuscular, subcutaneous injection).  
Treatment duration: 3 - 5 days.
- **Cattle, Buffaloes, Camels, Sheep, Goats, Pigs:** 0.15 - 0.40 ml per 10 kg B.wt. daily.
- Divide the dosage into separate injection sites to minimize the local tissue irritation.

## SAFETY PERIOD:

- For meat consumption: 14 days.
- For milk consumption: 5 days.

## REMARKS:

- Transient localized swelling may occur at the site of injection for few days.
- Not given with penicillins, calcium and iron preparations as interference action may occur.

**STORAGE:** Keep in dry, dark place below 30 °C.

**PACKING:** (50, 100) ml.