


# OXYDONE FORTE-30% LA

Injectable solution  
For livestock



**COMPOSITION:** Each ml contains:

**Oxytetracycline HCL**                      **300 mg**

## **PROPERTIES:**

- Gram-positive and Gram-negative pathogenic bacteria, certain Mycoplasma, rickettsiae, protozoa and Chlamydia. indicated in the treatment of a wide range of common systemic, respiratory, gastrointestinal, urogenital and local infections caused by, or associated with, organisms sensitive to oxytetracycline in cattle and pigs.
- Specific indications include:  
Calf diphtheria, foot abscess, mastitis, metritis, pasteurellosis and pneumonia, Erysipelas, mastitis-metritis-agalactia (MMA) syndrome, navel/joint ill.

**DOSAGE:** by deep intramuscular injection (single dose)

- **Cattle:** 1 ml per 15 kg, every 3-4 days
  - **Sheep, Goats:** 1 ml per 15 kg, every 3-4 days
  - **Pigs:** 1 ml per 15 kg body weight for 4-5 days
- For a prolonged duration of activity (4-5 days in cattle, 5-6 days in pigs) use 1 ml per 10 kg.

## **WITHDRAWAL PERIOD:**

**MILK:** Cattle, sheep, goats: 10 days.

**MEAT:** Cattle: 35 days, Sheep, goats, pigs: 28 days.

## **REMARK:**

Local transient may occur at the injection site.

## **PRECAUTION:**

- Not used in bobby calves
- During the period of tooth and bone development, including late pregnancy, may lead to tooth discoloration.

## **STORAGE:**

Keep in dry, dark place below 30 °C.

**PACKING:** 100, 50 ml.