


# OXYDONE – 500

(Bolus)



**COMPOSITION: Each Bolus contains:**  
**Oxytetracycline - HCL**

**500 mg**

## INDICATIONS:

- For sheep, goats, cattle, equines and poultry.
- Alimentary tract infections: enteritis, diarrhea, scours, bacillary dysentery.
- Respiratory tract infections: pneumonia, bronchitis, pleurisy, shipping fever
- Genital tract cases: metritis, vaginitis, prolapse.
- Calf diphtheria, septicemia, pyaemia, navel ill and joint ill infections.
- Skin infection, wounds, abscesses, foot rot, thrush, foul foot.
- Bacterial complications of viral infections.
- Control of post – operative and post parturient infections.
- Its used in poultry for blue comb disease paratyphoid loss of appetite associated with chronic respiratory diseases, hexamitiasis, infectious sinusitis of turkey

**DOSAGE:** given once daily, for 2-3 days. in severe cases, continue treatment for additional one day.

BODY WT. (KG)	SCOURES	RESPIRATORY DISEASES AND OTHER CASES.
CATTLE , CALVES: 45-90 135-255 225-450	1 2-3 3-4	2-4 4-10 8-20
SHEEP, GOATS, LAMBS, KIDS: UP TO 9.0 9.0 – 18.0 18.0 - 45.0	$\frac{1}{2}$ 1 1-2	$\frac{1}{2}$ - 1 1 1-2
EQUINE : UP TO 135 135 AND MORE	1-1 $\frac{1}{2}$ 2-3	- -

**POULTRY:** 1-2 tablets per 30 chickens.

## RAMARKS :

- If no improvement occurs after 36-48 hrs, re- diagnose the case
- If side – effects occur, discontinue treatment.

**SAFETY PERIOD:** 14 days for milk and meat consumption.

**STORAGE:** keep in dry, dark place below 30 °C.

**PACKING :** (50, 100 bolus)