

OXYTETRACYCLINE – 5%

For Bees

Water Soluble Powder

COMPOSITION: Each gm contains:

Oxytetracycline HCL **50 mg**

INDICATIONS:

For the control and treatment of American and European Foul Brood (*Spsigonia*) of bees.

DOSAGE AND APPLICATION:

▪ **Treatment:**

Dusting:

- Mix well (60) gm of the product with (240) gm of dry, thoroughly – milled and fine sugar. Then dust on the frames at a rate of (20) gm for each hive (2 gm per frame).
- Repeat treatment with an interval of (3) days between each application.

Sugar syrup:

- Dissolve well (6) gm of the product in small amount of water then mix thoroughly the solution with (1-2) litres of syrup (1:1 concentration) for each application.
- Repeat treatment (3-5) times (according to the severity of the case) with an interval of (3) days between each application.

- **Prevention:** Use half of the treatment dose for (3) times with an interval of (3) days between each application.

REMARKS:

- Apply medication in the evening, when all the foragers are inside the hive.
- Not to be applied during honey harvesting, otherwise, honey should be used for industrial purposes and not for human consumption.
- Medicate after honey – harvesting and stop treatment one month before the harvesting.
- Severely infected frames and boxes should be burnt.
- Dry and ventilate the hives.
- Use always clean and sterile frames and boxes.
- Old or weak queens should be changed.

PRECAUTION: Wash the hands after usage.

STORAGE: Keep in a dry, dark place below 30 °C.

PACKING: (60) GM.