

P-OXYRIDE 50%

**FUNGICIDE
WETTABLE POWDER**

**Class:
Multi-Site
Inorganic**

A protective fungicide with contact action. Effective in controlling many of the most fungal diseases on a wide range of agricultural crops .

COMPOSITION : Each kg contains:

COPPER OXYCHLORIDE 50% W/W (A.I)

PROPERTIES:

P-OXYRIDE 50% : A protective foliar fungicide with contact action, deposits must be on the crop before fungal spores begin to germinate.

P-OXYRIDE 50% : Control a wide range of fungal diseases on a wide range of Crops.

P-OXYRIDE 50% : Not phytotoxic for treated plants , but may cause injury to plum and peach foliage ,especially at low temperatures.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE GM/20L	APPLICATION TIME
Apricot Plum, Cherry, Peaches, Almond.	Shothole, Leaf Curl.	50	- Apply at bud swell but before and within one week of bud opening . - Repeat 7 - 10 days .
Apple, Pears	Scab	50	- Repeat 7 - 10 days.
Mango	Black Spot	50	- Apply every 4 weeks from early flowering to fruit set
Avocado	Anthracnose	50	Repeat At 28 Days.
Guavas	Anthracnose, Leaf Spot	50	-----
Banana	Leaf Spot	50	- Add 30 ml of summer oil. - Apply at 3 - 4 weekly intervals commencing december to may when weather conditions favour disease.
Vines	Downy Mildew	50	- First apply when shoots are 10 cm long, repeat at 10 - 14 day.
Citrus	Black Spot, Melanose, Smoky Blotch, Scab On Lemon.	40-60	- Add 30 ml of summer oil. (apply at 50 - 75% petal fall, use higher rate in coastal districts)
Potatoes	Early Blight, Late Blight	45-50	- Apply from crop emergence to crop maturity at 7 - 14 day .
Tomatoes	Early Blight, Late Blight, Bacterial Spot, Septoria Spot	45-50	- Seedbeds: apply each 7 days under wet weather conditions and repeat at 7 - 14 day .

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE GM/20L	APPLICATION TIME
Cucumber, Watermelon, Cucurbit, Squash	Bacterial Spot, Angular Leaf Spot	50	- Repeat 10 - 14 days.
Pepper	Leaf Spot, Leaf Canker	50	- Repeat every 7 days wet weather .
Cabbage, Cauliflower	Bacterial Spot, Downy mildew, Black Leaf Spot	50	- This product must not be applied to cabbage, when frost is likely to occur. - Repeat 10 - 14 days .
Lettuce	Bacterial Spot, Downy mildew, Anthracnose.	50	- This product must not be applied to lettuce, when frost is likely to occur. - Repeat 10 - 14 days .
Carrot	Early Blight	50	- Repeat 7 - 14 days .
Celery	Early Blight, Late Blight	50	- Repeat 7 - 14 days .
Onion	Downy Mildew	50	- Repeat 10 days .
Beans, Peas	Blights, Rust, Bacterial Spot, Chocolate Spot	50	- Repeat 10 - 14 days .
Tobacco	Wildfire, Angular Leaf Spot	80	- Repeat 7 days .
Coffee	Leaf Rust, Black Rot	38	---
	Blight	70	
	Leaf Spot	25	
Ornamentals	Fungal Leaf Spots, Downy mildew	60	- Repeat 7 - 14 days .

SAFETY PERIOD:

1 day on the above mentioned crops.

REMARKS:

Used only on above mentioned crops.

RE-ENTRY PERIOD:

Do not enter treated field before 24 hours at least after application.

COMPATIBILITY:

Compatible with most pesticides except alkaline materials.

• For more details about first aid & precautions please refer to first aid & precautions index.