

# PYLORA-24%

ACARICIDE / INSECTICIDE  
SUSPENSION CONCENTRATE

VAPCO

Class:  
Arylpyrrole

**Acaricide and Insecticide with mainly stomach and some contact action. Controls many of insects and mites species, including those resistant to Carbamate, Organophosphate and Pyrethroid insecticides and also chitin synthesis inhibitors.**

**COMPOSITION:** Each liter contains:

**CHLORFENAPYR**

**24% W/V (A.I)**

**PROPERTIES:**

**PYLORA- 24% :** Acaricide and insecticide with mainly stomach and some contact action. control of many species of insects and mites.

**PYLORA- 24% :** Applied on insects including those resistant to Carbamate, Organophosphate and Pyrethroid insecticides and also chitin synthesis inhibitors.

**PYLORA- 24% :** Has a very low residual effect.

**USES AND APPLICATION RATE:**

CROP	PEST	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Cotton	Tobacco Budworm	20-30	4
	Broad Mite	25	
	Two-spotted Red Spider Mite	20	
	Fall Armyworm		
Pepper, Eggplant, Cucumber, Okra Strawberry	Two-spotted Red Spider Mite, Rust Mite, Thrips, Cut Worms.	5-10	1
Muskmelon, Watermelon	Thrips	10-20	3
Cabbage	Diamondback Moth.	5-10	7
Tomato	Tomato Leafminer, Russet Mite Red Spider Mite .	5-10	3
Potato	Potato Tuber Moth	10-15	3
	Beetle		
	Thrips		
	Leafminer	15	

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PEST	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Apple	Apple Rust Mite, Leafminer, Two-spotted Red Spider Mite	5-15 20-30 10-20	7
Papaya	Broad Mite	6-10	2
Passion Fruit	Juno Longwing	6-10	3
Peach, Nectarine	Thrips, Leafminer	20-25	1
Grapes	Two-spotted Red Spider Mite	10-20	3
Citrus	Rust Mite, Scale Insect, Cut Worms	20-25	21
Onion	Thrips	5-10	7
Garlic	Thrips	10-20	3
Bean	Whitefly	20	3
	Thrips	10-15	
	Beetle		
Corn	Fall Armyworm	10-15	2
Ornamentals	Red Spider Mite, Cut Worms	5-10	---

#### CAUTION:

- Used only on the above mentioned crops with commitment with safety period.
- Don't use **PYLORA- 24% SC** on small seedling, newly transmitted to the field until 2 weeks.
- **PYLORA- 24% SC** may causes sensitivity if used on some ornamentals such as:  
Carnation, Dianthus, Kainchoe, Poinsettia, Roses, Salvia and Zinnia.

#### RE-ENTRY PERIOD:

48 hours from last application.

#### COMPATIBILITY:

Compatible with most other pesticides except bordeaux solution, mixing trials recommended before use.

• For more details about first aid & precautions please refer to first aid & precautions index.