

QUIZOVAP-10.8%

**HERBICIDE
EMULSIFIABLE CONCENTRATE**

VAPCO

**Class:
Aryloxyphenoxy-
Propionate**

Systemic herbicide, absorbed from the leaf surface, with translocation throughout the plant, moving in both the xylem and phloem, and accumulating in the meristematic tissue.

COMPOSITION : Each liter contains :

QUIZALOFOP-P-ETHYL

10.8% W/V (A.I)

PROPERTIES:

QUIZOVAP-10.8%: Systemic herbicide, absorbed from the leaf surface, with translocation throughout the plant, moving in both the xylem and phloem, and accumulating in the meristematic tissue.

QUIZOVAP-10.8%: Selective post-emergence to control of annual and perennial grass weeds in crops which are mentioned in the table.

QUIZOVAP-10.8%: Tolerant to most non-graminaceous crops.

USES AND APPLICATION RATES:

CROPS	WEEDS	WEED STAGE	APPLICATION RATES	SAFETY PERIOD	NOTICE
Beetroot Cabbage Carrots Cauliflower Chickpeas Clover Cucumber Faba Beans Peas Melon Onions Potatoes Pumpkins Radish Tomatoes Soybeans Sunflower	Annual Ryegrass, Brome Grasses	3 leaf to early tillering	23-35 ml / downm	14 days 9 weeks 10 weeks 14 days 12 weeks -----	Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering.
	Volunteer Barley, Volunteer Wheat	3 leaf to early tillering	23 ml / downm	14 days 12 weeks 9 weeks 9 weeks 18 weeks 10 weeks	Apply when weeds are actively growing
	Wild Oats	3 leaf to mid tillering	11.5-23 ml / downm	9 weeks 21 days 4 weeks 12 weeks 9 weeks	Apply when weeds are actively growing. Use a lower rate only under light weed pressure and/or when weeds are at the 1-3 leaf stage. Use higher rate under heavy weed pressure and/or when weeds are tillering.
	Couch Grass	2-4 leaf seedling	46 ml / downm		Apply when weeds are actively growing. This rate will only control seedling couch grass which is less than 2 months old and established from seed.
	Johnson Grass	2 leaf to 3 tiller	46-69 ml / downm		Apply when weeds are actively growing. Use higher rate under heavy weed pressure, and/or when weeds have commenced tillering.

Donum = 1000 m²

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

REMARKS:

- Calibration of sprayer is important before application.
- Its favorable to add a sticking agent **SUPER SURFIX** (VAPCO'S product) before spraying.
- Do not apply **QUIZOVAP-10.8% EC** before the five true leaf stage of the crop.
- Do not apply to crops that are stressed by any cause (such as severe cold weather, disease, drought or waterlogging).
- Do not apply to weeds that are stressed by any cause (such as weather conditions, drought, waterlogging) or not actively growing.
- Grass weeds under stress frequently become less susceptible to herbicide activity. This may result in an incomplete kill or suppression of weeds only.
- Do not apply under conditions of high temperature and low humidity.
- Do not apply if rainfall is expected within 3 hours.
- Do not plant cereal crops into the treated area for a period of 18 weeks after application of **QUIZOVAP-10.8% EC**.
- Use separate sprayer for herbicides or clean the used sprayer after application.

RE-ENTRY PERIOD:

24 hours after the last application.

GRAZING RESTRICTIONS:

Don't graze or feed animals from treated area for at least 14 days for Clover and Peas, other crops 12 weeks.

COMPATIBILITY:

Can be used in combination with post-emergence broad-leaved herbicides.

• For more details about first aid & precautions please refer to first aid & precautions index.