

RATRYL-0.75%

RODENTICIDE
TRACKING POWDER

VAPCO

Class:
Coumarin
anticoagulant

Is an anticoagulant rodenticide which has shown exceptional activity against all rodent species, and can be used in all places where rodents can live.

COMPOSITION: Each kg contains:

COUMATETRALYL

0.75% W/W (A.I)

PROPERTIES:

RATRYL-0.75%: Gives excellent control of rats and house mice. A daily dosage ingested several times.

RATRYL-0.75%: Is readily accepted and do not cause any development of poison shyness.

RATRYL-0.75%: Causes inconspicuous and painless death of rats, just as if they had died of senility. There are no visible symptoms, so that other rats in the colony are not warned.

RATRYL-0.75%: Is no danger of secondary poisoning to cats.

USES AND RECOMMENDATIONS:

- **RATRYL-0.75%:** Is recommended to control rats in cellars, on farms, in stables housing large and small animals, in zoological gardens, slaughter-houses, cattle yards, goods depots and terminals, refuse tips, sewers, sewage works, port installations, on ships, in coal mines....etc.
- **RATRYL-0.75%:** Is highly recommended for use in area-wide rat eradication programmes in cities, towns and rural districts. Control operations must be preceded by surveys to determine rat infestation density, and followed up to regular long-term inspections and re-treatments to maintain the achieved state of clearance.
- **RATRYL-0.75%:** Product may used in control rats in agricultural crops.

METHOD OF APPLICATION: May be used with two methods:

- **RATRYL-0.75%** is ready for use product used to control rats. It applied by sprinkling it in a layer, in hiding places and on rat runways. The deposit must be regularly checked and replenished if necessary.
- **RATRYL-0.75%** can be mixed with a wide range of cereals (Corn, Wheat, Barley ... etc) to make a dry bait formulation. Users can prepare baits themselves by mixing 1 part of **RATRYL-0.75%** with 19 parts of bait material.
The following mixture show exceptional effectiveness: 550 gm of ground Maize + 350 gm rolled Oats + 50 gm of castor Sugar + 50 gm of **RATRYL-0.75%**.

• For more details about first aid & precautions please refer to first aid & precautions index.

REMARKS:

- Do not touch **RATRYL-0.75%** with bare hands as human scent is a warning to rodents.
- In case of using a tracking powder, sprinkle the powder around rat hole and runways until signs of rat and mouse activity decrease.
- Do not place powder in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.
- Powder not touched after 5-8 days should be moved to a new Location.
- Maintain an uninterrupted supply of fresh powder for 15 days or until all sign of feeding touching has disappeared.
- Neighbors and children should be warned that rodenticide has been applied.
- Remove all dead rats by placing them in tight-closing bags, after dusted with insecticide. The carceases should be handled with rubber or polyethylene gloves and all of the dead rats either burned or buried.

• For more details about first aid & precautions please refer to first aid & precautions index.