

ROVER-50%

**FUNGICIDE
WETTABLE POWDER**

VAPCO

Class:
Dicarboximide

Protective and curative fungicide acts by contact action, control many fungal diseases on a wide range of crops.

COMPOSITION: Each kg contains:

IPRODIONE

50 % W/W (A.I)

PROPERTIES:

ROVER-50%: Contact fungicide with protective and curative action, inhibits germination of spores and growth of fungal mycelium.

ROVER-50%: Control of Rots, Blight and Shot hole on a wide range of crops.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE GM/ 20 LT. WATER	SAFETY PERIOD (DAYS)
Stone fruits: Peach, Apricot, Cherry, Nectarine, Plum and Prunes.	Blight, Brown rot, Shot hole, Leaf spot, Fruit rot, Scab	20 – 30	15
Strawberries	Gray mold, Stem rot, Soft rot, Anthracnose, Stem canker	20 – 30	3
Grapes	Bunch Rot	20 – 30	7
Almonds	Brown Rot, Blight, Shot Hole	22	---
Caneberries	Botrytis Fruit Rot	22 - 44	0
Peanuts	Sclerotinia Blight	44	10
Potatoes	Early blight, White rot	20 – 30	14
Carrots	Alternaria blight, black crown rot	22 - 44	0
Lettuce	Lettuce drop, bottom rot	33 - 44	14
Broccoli	Black leg	44	0
Chinese mustard	Alternaria leaf spot	22	10
Beans (Snap, Dry, Lima)	Gray Mold, White Mold	26 - 44	14
Garlic (One Application / Season)	Sclerotinia Rot	20 – 30	2
Dry bulb onion	Alternaria Leaf Spot	22	10

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE GM/ 20 LT. WATER	SAFETY PERIOD (DAYS)
Canola	Sclerotinia Stem Rot	20 - 30	---
Turf	Dollar Spot	90	21
	Brown Patch	120 – 250	
	Leaf Spot	120	
Ornamentals	Botrytis spp	20	---

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

48 hours after the last application.

COMPATIBILITY:

Compatible with most pesticides, it's preferable to make compatibility test before use.

• For more details about first aid & precautions please refer to first aid & precautions index.