


# RENAMIDE-5%

Injectable Solution  
Diuretic



**COMPOSITION :** Each ml contains :  
**Furosemide 50 mg**

## PROPERTIES :

- Potent diuretic with rapid onset action
- Increased Na and water excretion without loss of potassium.
- Effective even in cases of glomerular – infiltration impairment .

## INDICATIONS :

- For cattle , equines, dogs, cats, pigs.
- Pathological oedema of the udder and perineum of the cow and mare at time of parturition .
- Preputial or scrotal oedema of male animals .
- Supportive treatment in the elimination of fluids in cases of ascites, hydrothorax , and bronchopneumonia , hydropericardium , from joints and tendon sheaths of bursae .
- Supportive treatment in laminitis, paralytic myoglobinuria of the horse .
- Protein deficiency oedema due to renal proteinuria , liver lesions caused by protein metabolism disorders, parasitic infestations or mal- nutrition seen as generalised skin oedema and ascites .
- Skin oedema due to blood vessels lesions of toxicity, trauma, allergy , inflammations , wounds, surgery.

## DOSAGE : ( By Intramuscular Or Slow Intravenous Injection )

- **Equines :** 5-10 ml once or twice daily with intervals of 6-8 hours.
- **Cattle :** 10 ml one time, or 5 ml twice with interval of 6-8 hours .
- **Dogs ,Cats:** 0.25-0.5 ml /5 kg b.wt. once or twice daily with interval of 6-8 hours.
- **Pigs :** 1 ml/10 kg b.wt once or to be determined empirically.

**SAFETY PERIOD :** For meat and milk consumption : 2 days .

## REMARKS :

- ◀ Not used in cases of renal failure with anuria, oliguria, acute glomerular nephritis , electrolytes deficiency disease or over dosage with digitalis .
- ◀ The therapeutic effect may be impaired by increased water intake .
- ◀ Diuresis may be enhanced with oral NaCl intake.
- ◀ In hypocalcaemic animals , tetany may occur.

## STORAGE :

Keep in dry, dark place below 30 °C

**PACKING :** (50,100) ML .