

# SELFAT-53.5%

**FUNGICIDE  
WETTABLE POWDER**

**VAPCO**

Class:  
**Inorganic compound<sup>(1)</sup>  
Dithiocarbamate<sup>(2)</sup>**

**Foliar fungicide with contact, preventive action. Used to control many fungal diseases such as: Downy mildew, Blights, Leaf Spots and Scabs.**

**COMPOSITION:** Each kg contains:

**COPPER OXYCHLORIDE<sup>(1)</sup>**

**37.5 % W/W (A.I)**

**ZINEB<sup>(2)</sup>**

**16 % W/W (A.I)**

## **PROPERTIES:**

**SELFAT-53.5%:** Foliar fungicide with contact, preventive action. Deposits must be on the crop before fungal spores begin to germinate.

**SELFAT-53.5%:** Used to control many fungal diseases such as: Downy mildew, Blights, Leaf Spot and Scab.

**SELFAT-53.5%:** Non-phytotoxic, except for Zinc-sensitive plants such as Tobacco and Cucurbits, and may cause injury to Plum and Peach foliage, especially at low temperatures.

## **USES AND APPLICATION RATE:**

<b>CROP</b>	<b>DISEASE</b>	<b>APPLICATION RATE GM/ 20 LT. WATER</b>	<b>SAFETY PERIOD (DAYS)</b>
<b>Squash, Cucumber, Pumpkins, Watermelon</b>	Anthracoese, Downy mildew	40 – 60	3
<b>Potatoes</b>	Late blight, Alternaria Spp., Anthracoese , Downy mildew	50	7
<b>Tomatoes</b>	Late blight, Downy mildew	40 – 60	7
<b>Solanaceous (Eggplant, Sweet Peppers, Chilli Peppers (But Not Black Pepper)</b>	Anthracoese , Downy mildew	40 – 60	3
<b>Brassica Vegetables (Cabbage, Brussels, Sprouts Cauliflower, Radish, Broccoli)</b>	Alternaria Spp., Downy mildew	40 – 60	28
<b>Garlic, Onion, Shallot, Aromatic Herbs and Hops.</b>	Downy mildew.	40 – 60	15
<b>Lettuce , Leek</b>	Downy mildew	40 – 60	21
<b>Carrot</b>	Alternaria Spp., Downy mildew	40 – 60	15
<b>Celery</b>	Downy mildew , Septoria leaf spot	40 – 60	21

• For more details about first aid & precautions please refer to first aid & precautions index.

<b>CROP</b>	<b>DISEASE</b>	<b>APPLICATION RATE GM/ 20 LT. WATER</b>	<b>SAFETY PERIOD (DAYS)</b>
<b>Fruit trees</b>	Scab, Leaf spot	60 – 80	15
<b>Citrus Tree</b>	Foot rot, Alternaria Spp., Citrus Bacteriosis, Sooty mold.	40 – 60	15
<b>Peaches, Nectarines, Plums, Apricots, Cherries</b>	Cankers, Brown rot.	40 – 60	28
<b>Pomaceous Trees (Apple, Pear, Quince Trees)</b>	Brown rot , Scab	40 – 60	28
<b>Vine Trees</b>	Anthrachnose , Downy mildew	40 – 60	15
<b>Chickpea</b>	Ascochyta Leaf spot	40 – 60	15
<b>Green Pea</b>	Anthrachnose, Downy mildew	40 – 60	21
<b>Grain Beans</b>	Alternaria Spp., Anthracnose, Bean rust	40 – 60	21
<b>Green Beans</b>	Anthrachnose , Downy mildew	40 – 60	21
<b>Sugar Beet</b>	Leaf spot	4 – 6 kg / ha	28
<b>Olive Tree</b>	Olive Peacock spot	40 – 60	15

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

**RE-ENTRY PERIOD:**

48 hours after the last application.

**COMPATIBILITY:**

Compatible with most common pesticides, but it is favorable to make compatibility test before use.

• For more details about first aid & precautions please refer to first aid & precautions index.