

SPIDERON-10%

ACARICIDE
WETTABLE POWDER

VAPCO

Class:
Mite Growth Inhibitor

Non systemic acaricide used to control many species of mite with contact action, and long residual activity .

COMPOSITION: Each Kg contains:

HEXYTHIAZOX

10% W/W (A.I)

PROPERTIES:

SPIDERON-10% : Non systemic acaricide with contact action.

SPIDERON-10% : Highly effective in controlling immature motile stages of target mite species that are sprayed or move onto treated surfaces

SPIDERON-10% : Provide extended residual control of pest mite species without disrupting beneficial insects .

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATES GM/20 LT. WATER	SAFETY PERIOD (DAYS)
Citrus	European Red Mite.	8 - 10	7
Apple, Pears	European Red Mite, Two-spotted Spider Mite.	8 - 10	7
Peaches, Apricots, Plums, Cherries, Nectarines	European Red Mite.	8 - 10	7
Grapes	European Red Mite.	8 - 10	7
Strawberries	European Red Mite.	8 - 10	1
Cucumbers, Squash, Melons, Pumpkins	Two-spotted Spider Mite.	8 - 10	7
Tomatoes, Eggplant, Peppers, Peas	Two-spotted Spider Mite.	8 - 10	1
Beans	European Red Mite.	8 - 10	14
Corn	Two-spotted Spider Mite.	8 - 10	14
Cotton	Two-spotted Spider Mite.	8 - 10	14
Figs	Two-spotted Spider Mite.	8 - 10	1
Ornamentals	European Red Mite.	8 - 10	-----

Caution:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

Do not enter treated field before 24 hours at least after applications.

COMPATIBILITY:

Compatible with most common pesticides, use the solution direct after mixing and it's preferable to make test before spraying .

• For more details about first aid & precautions please refer to first aid & precautions index.