

SUPERTAK-10%

**INSECTICIDE
EMULSIFIABLE CONCENTRATE**

VAPCO

Class:
Pyrethroid

Pyrethroidal, highly effective insecticide which controls a wide range of pests attack agricultural crops.

COMPOSITION: Each liter contains:

ALPHA-CYPERMETHRIN

10% W/V (A.I)

PROPERTIES:

SUPERTAK-10%: Pyrethroidal, highly effective insecticide which controls a wide range of pests attack agricultural crops.

SUPERTAK-10%: Economic pesticide effective at a lower dose.

SUPERTAK-10%: Non-phytotoxic if used as recommended.

SUPERTAK-10%: Non-systemic insecticide acts as contact and stomach poison.

SUPERTAK-10%: Control insects which become resistant to other insecticides.

SUPERTAK-10%: Can be used as protective spray since it has a long residual effect.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Tomato, Lettuce, Cucumber, Bean, Peas. Cabbage, Potato	Cutworms, American Boll Worms,	3.5 - 6	3
	Leafworm, Beetles, Whitefly, Leafminer, Scale insects, Aphid		14
Apple, Pear, Peach, Cherries, Apricot, Almond, Grape.	Scale insects, Cutworm, Beetles, Leafminer, Aphid, Thrips, Whitefly	6	7
Tobacco, Corn, Sunflower	Native Bud Worm, Corn Armyworm	6	7
Cotton	Cotton Boll Worm, Native Bud Worm	6	7
Coffee	White Coffee Leafminer	6	7
Tea	Tea Tortrix	6	9
Rice	Common Armyworm	4	7
Oil Palm	Bag Worm, Caterpillars	2	7
Asparagus	Garden Weevil	20	1

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest

RE-ENTRY PERIOD:

48 hours after the last application.

COMPATIBILITY:

Compatible with most other insecticides and fungicides except highly acidic and alkaline compounds.

*** OTHER FORNULATION & TRADE NAME:**

SUPERTAK – 5% EC, SUPERKALL – 25% EC.

• For more details about first aid & precautions please refer to first aid & precautions index.