

TEBIZOLE-25%

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Triazole

Systemic fungicide with protective and curative action to control many fungal diseases on a wide range of crops and to protect seed during germination stage.

COMPOSITION: Each kg contains:

TEBUCONAZOLE

25 % W/W (A.I)

PROPERTIES:

TEBIZOLE-25%: Systemic fungicide with protective and curative action.

TEBIZOLE-25%: Fungicide recommended to control diseases such as Rust, Powdery mildew, Early leaf spot, Late leaf spot, Net blotch, Stem and pod rot, Rhizoctonia wilt, Monilia, Scaled, Scab, Septoria spot, Early blight, Stem canker on Almond, Eggplant, Barley, Bean, Cucumber, Grapes, Peanut, Plum, Potatoes, Sunflower, Sweet pepper, Tomatoes, Wheat, Cereal and Corn.

TEBIZOLE-25%: Effective against various Smut and Bunt diseases of Cereals.

TEBIZOLE-25%: Slightly toxic to Mammals, toxic to Fish and Bees

USES AND APPLICATION RATE:

A.FOLIAR APPLICATION:

CROP	DISEASE	APPLICATION RATES GM / HA	SITE	NO. OF APPLICATIONS	SAFETY PERIOD (DAYS)
Tomatoes	Early blight	600	field & green house	1-3	14
Eggplant	Powdery mildew	600	field & green house	3	14
Cucurbits (Melons, Cucumbers, Squash, Pumpkin)	Powdery mildew, Fruit blotch, Downy mildew	460 – 690	field & green house	4 – 6	10 – 14
	Gummy stem blight (Melons, Squash and Pumpkin only)	920		3	
Lettuce	Sclerotia Rot	600	field	2	35
Sweet Pepper	Powdery mildew	1 kg	field	1-2	7
Onion	Purple blotch	750	field	1	21
Almond	Monilia, Powdery mildew	200	field	1	21
Plum	Monilia, Powdery mildew	200	field	1	21

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATES GM / HA	SITE	NO. OF APPLICATIONS	SAFETY PERIOD (DAYS)
Grapes	Powdery mildew	600	field	2	14
Banana	Leaf spot	390	field	6	14
Papaya	Black spot	800	field	6	7
Peanut	Early leaf spot, Late leaf spot, Rust, Net blotch	1 kg	field	1	21
French Bean, Kidney Bean	Leaf spot, Rust	1 kg	field	1 – 2	3
Sunflower	Stem canker, Stem rot	500	field	1	21
Barley	Rust, Powdery mildew, Head scab, Septoria spot.	750	field	1	30
Wheat soft	Rust, Powdery mildew, Scald, scab	500	field	1-2	60
Wheat	Septoria spot	1 kg	field	1-2	60
Rice	Leaf and Ear blight, Leaf scald	1 – 1.5 kg	field	1	35
	Rice blast	1 kg			
Roses	Black spot, Powdery mildew, Rust	3.4 kg	field & green house	3	14
Flower	Leaf spot, Powdery mildew, Rust, Southern blight	3.4 kg	field & green house	3	14
Turf	Dollar spot, Copper spot, Powdery mildew, Rusts	8 kg	On Golf Course Turf Only	6	28

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

• For more details about first aid & precautions please refer to first aid & precautions index.

B. SEED TREATMENT:

TEBIZOLE-25% used to protect seeds during germination as follow:

CROPS	DISEASE	APPLICATION RATES GM/100 KG SEED
Barley	Barley leaf stripe, Covered smut, False loose smut, Seed rot and Seedling blight caused by <i>Fusarium</i> , True loose smut.	4 – 12
Oats	Loose smut.	4 – 12
Wheat	Common bunt (stinking smut), Loose smut, Seed rot and Seedling blight caused by <i>Fusarium</i> . Suppression of Common root rot caused by <i>Cochliobolus sativus</i> .	4 – 12
Corn	Head smut, Loose smut.	30

WARNING:

Treated seeds are not suitable for human consumption.

TYPE OF APPLICATION:

Use the recommended dose of **TEBIZOLE-25%** with 100 kg seeds and mix with 300-500 ml of water to obtain slurry mixture.

RE-ENTRY PERIOD:

12 hours after the last application.

COMPATIBILITY:

Compatible with most common used pesticides, but it's recommended to make compatibility test before use.

• For more details about first aid & precautions please refer to first aid & precautions index.