

TETRAMISOLE – 30

Water Soluble Powder

The Safe & Broad Spectrum Vermicidal, Larvicidal Anthelmintic For Livestock



COMPOSITION : Each gm contains :

Tetramisole HCL 300 mg

PROPERTIES :

- Tetramisole is broad spectrum and powerful anthelmintic (imidothiazole group) against nematodes (round worms) of the gastrointestinal tract and the lung worms of ruminant .
- Tetramisole at the therapeutic dose . is safe even for the pregnant animals
- Tetramisole exert its anthelmintics action by paralyzing the nematode .
The paralysis id due to sustained muscle contraction which leads to neuro – muscular paralysis of the parasites.

INDICATIONS:

For the treatment and control of parasitic gastro – enteritis and verminous bronchitis caused by round worms (Nematodes):

- **Ruminants, Rabbits:**
 - **Gastro – intestinal worms:** Ascaris, Nematodirus, Haemonchus, Ostertagia, Cooperia, Trichostrongylus.
 - **Lung worms:** Dictyocaulus.
- **Poultry:** Ascaridia, Heterakis, Capillaria, Amidostomum.

DOSAGE:

- Given with water or feed.
- Give single dose each time.
 - **Sheep, goats, cattle, camels:** 0.5 gm per 10 kg B. wt.
(15 g active sub. / kg body wt.).
Repeat for prevention every 4-6 months.
 - **Poultry, Pigeons:** 0.06 gm per kg B. wt.
(20 mg active sub. / kg body wt.)
Repeat every 3 months.
 - **Rabbits:** 0.06 gm per kg B. wt. (20 mg active sub. / kg B. wt.)
Repeat twice per year.

SAFETY PERIOD:

- For meat and eggs consumption: 7 days.
- For milk consumption: 1 day.

REMARKS:

- Organophosphates insecticides are toxic if applied within (14) days of treatment.
- Avoid treatment of debilitated or sick animals.
- Occasionally, salivation , slight diarrhea and coughing may occur in some animals.

STORAGE: Keep in a dry, dark place below 30 °C.

PACKING: (100, 250, 500, 1000) GM.