TITANOL-25%

FUNGICIDE WETTABLE POWDER



Foliar fungicide, with protective and curative action, used to control many fungal diseases on a wide range of agricultural crops.

COMPOSITION: Each kg contains:

BITERTANOL 25 % W/W (A.I)

PROPERTIES:

TITANOL-25%: Foliar fungicide with protective and curative action, inhibits spore germination

and prevent mycelium development.

TITANOL-25%: Used to control many fungal diseases such as: Scab, Monilia, Powdery Mildew,

Spots on a wide range of agricultural crops.

TITANOL-25%: Slightly toxic to mammals, not toxic to bees but toxic to fish.

USES AND APPLICATION RATE:

| CROP | DISEASE | APPLICATION RATE GM/ 20 LT. WATER | SAFETY PERIOD (DAYS) |
|---|----------------------------|--------------------------------------|-------------------------|
| Apple, Pears | Scab, Powdery mildew. | 12 - 20 | 21 |
| Cherry, Plum, Peach, Apricots, Nectarine | Powdery mildew. | 12 - 16 | 21 |
| | Monilia | 20 - 24 | |
| | Canker | 20 | |
| Cucumber, Squash | Powdery mildew. | 12 - 20 | 14 |
| Sugarbeet | Powdery mildew, Leaf spot. | 12 - 20 | 30 |
| Banana | Mycosphaerella | 5 | 14 |
| Roses and Ornamentals | Powdery mildew, Rust. | 16 - 32 | |
| Forest Trees | Rust, leaf blotch | 20 - 40 | |

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest,

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY:

Compatible with most pesticides.

[•] For more details about first aid & precautions please refer to first aid & precautions index.