

TRIVAP-75%

INSECTICIDE
(INSECT GROWTH REGULATOR IGR)
WETTABLE POWDER

VAPCO

Class:
Triazine

An insect growth regulator with contact action, which interferes with moulting and pupation. When used on plants, it acts as systemic insecticide.

COMPOSITION: Each kg contains:

CYROMAZINE

75% W/W (A.I)

PROPERTIES:

TRIVAP-75%: Is an insect growth regulator with contact action, which interferes with moulting and pupation. When used on plants, it acts as systemic insecticide; when applied to the leaves it exhibits a strong translaminar effect, but when applied to the soil it is taken up by roots and translocated acropetally.

TRIVAP-75%: Does not kills adult insect nut it limits reproduction through weakening egg hatching and partial sterility of the eggs.

TRIVAP-75%: Used as a foliar spray to control Leafminer on vegetables such as: Celery, Melons, Tomatoes, Lettuce, Eggplant, Potatoes, Mushrooms and Ornamentals.

USES AND APPLICATION RATE:

CROP	PEST	APPLICATION RATE GM / 20 LT. WATER	SAFETY PERIOD
Open field: Squash, Pumpkin, Bean, Peas, Muskmelon, Watermelon, Carrots, Onion, Eggplant, Potato, Parsley, Lettuce, Pepper, Tomatoes.	Leafminer	25 – 35 Start spraying when the first infestation appears with 50 lt./donum and re-spray every 7-10 days if necessary.	3 days for: Tomatoes, Cucumber, Eggplant, Muskmelon, Watermelon, Squash and Pumpkin.
Inside green houses: Tomato, Cucumber, Pepper, Eggplant, Strawberries, Ornamentals.			14 days for: Bean, Peas, Carrots, Lettuce, Pepper, Potato and Onion.

CAUTION:

Donum = 1000 m²

Used only on above mentioned crops and follow the recommended safety period before harvest.

REMARKS:

Do not apply this product through any type of irrigation system.

RE-ENTRY PERIOD:

24 hours after the last application.

COMPATIBILITY:

Compatible with most known pesticides, but mixing with alkaline compound such as: Lime sulfur and Copper compounds may reduce effectivity.

• For more details about first aid & precautions please refer to first aid & precautions index.