



# TYLOSIN-100%

Water soluble Powder  
For Poultry, Turkeys, Pigeons

## COMPOSITION: Each 113gm contains:

Tylosin (As Tylosin Tartrate)                      100 gm

## PROPERTIES:

- Tylosin as a strong macrolide antibiotic covers a wide spectrum of Gm+ve bacteria such as: Staph, Strept, Diplococcus, Coryne, Clostridia and other bacterial infections secondary to viral diseases.
- Very effective against mycoplasma infections and affects also some rickettsias (Group psittacosis – chlamydiosis ) and Hemophilus gallisepticum.
- Some Gm-ve bacteria ( Pasteurella, Campylobacter ) also are affected by Tylosin.

## INDICATIONS:

- For prevention and treatment of: Mycoplasmosis (CRD), Air succulitis, Infectious synovitis, Infectious sinusitis, Chlamydiosis, Spirochaetosis.
- Stress conditions due to vaccination, debeaking, transporting, rehousing, changes of feed and weather.

## DOSAGE:

- **Solution preparation:** Add 4 liters of water to Tylosin 100% (not vice versa) and dissolve well then add to the water tank.

### Treatment:

- Respiratory or general infections in poultry: (0.05% solution) 100 gm (activity) per 200 litres of water for 3-5 days according to severity of the disease.  
Spirochaetosis: (0.5% solution) used as preventive and for control.
- Pigeons: 0.015 – 0.025 gm / kg b. wt. 4 times a day.  
(15-25 mg activity / kg B.wt. / 4 times daily).

- **Prevention program:**

Use 100 gm (activity) per 200 liters of water as follows:

- **Broilers:** 1<sup>st</sup> week of age for 3 days.  
4<sup>th</sup> week of age for 1 day.
- **Pullets:** 1<sup>st</sup> week of age for 3 days.  
4<sup>th</sup> week of age for 1 day.



9<sup>th</sup> and 20<sup>th</sup> week of age for 2 days.

- **Parent stocks:** 1<sup>st</sup> week of age for 5 days.  
4<sup>th</sup>, 9<sup>th</sup>, 16<sup>th</sup>, 20<sup>th</sup>, and 24<sup>th</sup> week of age for 2 days.
- **Turkeys:** 1<sup>st</sup> week of age for 5 days.  
3<sup>rd</sup> week of age for 1 day.

**WITHDRAWAL PERIOD:** For meat consumption: 7 days.

**REMARK:** Not recommended for table-eggs layers.  
During treatment, eggs not used for human consumption.

**STORAGE:** Keep in dry, dark place below 30 °C.

**PACKING:** 113 gm and others.